

AD-A142 136


THEORETICAL METHODS IN THE CALCULATION OF THE
BREMSSTRAHLUNG RECOIL FORCE IN A NONEQUILIBRIUM
BEAM-PLASMA SYSTEM(U) HARRY DIAMOND LABS ADELPHI MD
H E BRANDT MAY 84 HDL-TR-2009


1/1

UNCLASSIFIED

F/G 12/1


NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A142 136



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM												
1. REPORT NUMBER HDL-TR-2009	2. GOVT ACCESSION NO. AD-4142136	3. RECIPIENT'S CATALOG NUMBER												
4. TITLE (and Subtitle) Theoretical Methods in the Calculation of the Bremsstrahlung Recoil Force in a Nonequilibrium Beam-Plasma System	5. TYPE OF REPORT & PERIOD COVERED Technical Report													
7. AUTHOR(s) Howard E. Brandt	6. PERFORMING ORG. REPORT NUMBER													
9. PERFORMING ORGANIZATION NAME AND ADDRESS Harry Diamond Laboratories 2800 Powder Mill Road Adelphi, MD 20783	8. CONTRACT OR GRANT NUMBER(s)													
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Materiel Development and Readiness Command Alexandria, VA 22333	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Program Ele: 61101A													
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE May 1984													
	13. NUMBER OF PAGES 48													
	15. SECURITY CLASS. (of this report) UNCLASSIFIED													
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE													
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.														
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)														
18. SUPPLEMENTARY NOTES HDL Project: A10225 DRCMS Code: 61110191A0011 DA Project: 1L161101A91A														
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <table border="0"> <tr> <td>Radiation</td> <td>Plasma physics</td> <td>Bremsstrahlung</td> </tr> <tr> <td>Microwaves</td> <td>Relativistic electron beams</td> <td>Beam-plasma systems</td> </tr> <tr> <td>Plasma turbulence</td> <td>Radiative instability</td> <td>Nonlinear plasma physics</td> </tr> <tr> <td>Relativistic plasma</td> <td></td> <td></td> </tr> </table>			Radiation	Plasma physics	Bremsstrahlung	Microwaves	Relativistic electron beams	Beam-plasma systems	Plasma turbulence	Radiative instability	Nonlinear plasma physics	Relativistic plasma		
Radiation	Plasma physics	Bremsstrahlung												
Microwaves	Relativistic electron beams	Beam-plasma systems												
Plasma turbulence	Radiative instability	Nonlinear plasma physics												
Relativistic plasma														
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <p>The reduction of the lowest order nonlinear bremsstrahlung recoil force on the bare charge of an unperturbed relativistic test particle in a nonequilibrium beam-plasma system is derived from first principles. The purpose is to identify many of the important physical assumptions and mathematical techniques involved in Tsytovich's theory of nonlinear bremsstrahlung and radiative instability.</p>														

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

1

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CONTENTS

	<u>Page</u>
1. INTRODUCTION	5
2. REDUCTION OF A COMPONENT OF THE BREMSSTRAHLUNG RECOIL FORCE	8
3. CONCLUSION	37
LITERATURE CITED	38
SELECTED BIBLIOGRAPHY	39
DISTRIBUTION	41

Accession For

NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	

Classification/
Availability Codes
and number
of copies

A-1



1. INTRODUCTION

For a nonequilibrium beam-plasma system, the lowest-order Born approximation for the nonlinear force $\vec{F}_\alpha^{(1)}$ on the bare charge of an unperturbed relativistic test particle is given by¹⁻³

$$\vec{F}_\alpha^{(1)} = \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int d\vec{k} \vec{k} \frac{\vec{v}_\alpha \cdot \vec{E}_\vec{k}}{\omega + i\delta} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) . \quad (1)$$

Here α is a label designating the species of the test particle of charge e_α and velocity \vec{v}_α , t is the time, \vec{k} is the wave vector, and ω is the frequency corresponding to the Fourier component $\vec{E}_\vec{k}$ of the total field, $\delta(x)$ is the one-dimensional Dirac delta function, δ is a small imaginary part, and $d\vec{k} \equiv d^3\vec{k} d\omega$. Equation (1) differs from equation (7) of Akopyan and Tsytovich¹ by an additional factor of $(2\pi)^{-3}$ appearing in the latter because of the different Fourier transform convention chosen there.² If the part of the contribution to the force, equation (1), which is associated with recoil of the unperturbed bare test particle due to bremsstrahlung, is averaged over the random phase of the bremsstrahlung field, it reduces to

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle = \langle \vec{F}_\alpha^{\text{grad}(1a)} \rangle + \langle \vec{F}_\alpha^{\text{grad}(1b)} \rangle , \quad (2)$$

where

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1a)} \rangle &= \frac{1}{16\pi^2} e_\alpha^2 e_\beta^2 \int \frac{d^3\vec{p}_\beta d\omega d^3\vec{k} d^3\vec{k}'}{(2\pi)^3} \frac{\vec{k}}{\omega^2 (\vec{k} \cdot \vec{v}_\alpha)^4} \\ &\times |E_\vec{k}^{\sigma(0)}|^2 |e_{kj}^\sigma \Lambda_{ij}^{(\beta)}(\vec{k}, -\vec{k}) G_{ip}(-\vec{k}) v_{\alpha p}|^2 \\ &\times (\vec{k} - \vec{k}') \cdot \vec{v}_\beta f_{p\beta}^{R(0)} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha + (\vec{k} - \vec{k}') \cdot \vec{v}_\beta) \end{aligned} \quad (3)$$

is defined and calculated in this report, and $\langle \vec{F}_\alpha^{\text{grad}(1b)} \rangle$ is defined by equations (25), (26), and (47) below but calculated in another report.* The quantity $\langle \vec{F}_\alpha^{\text{grad}(1b)} \rangle$ is not mentioned by Akopyan and Tsytovich,¹ but it is nonvanishing and makes an important contribution to nonlinear bremsstrahlung.

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

²H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second-Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1995 (August 1983).

³H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1994 (September 1983).

*H. E. Brandt, *Collective Bremsstrahlung Recoil Force on the Bare Charge of an Unperturbed Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, 1983, (to be published).

The quantity $\langle \vec{F}_\alpha^{\text{grad}}(1a) \rangle$ given by equation (3) corresponds to equation (27) of Akopyan and Tsytovich.¹ Their equation (27) is expressed in Gaussian units, whereas equation (3) above is in MKS units. This results in an additional overall factor of $(4\pi)^2$ in Akopyan and Tsytovich.¹ Also there is an additional factor of $(2\pi)^{-6}$ in their work because of different Fourier transform conventions. However, their equation (27) also differs from equation (3) here in that they omit an overall factor of 2. Also, the arguments k and $-\kappa$ of $\Lambda_{ij}^{(\beta)}(k, -\kappa)$ in equation (3) appear incorrectly interchanged in equation (27) of Akopyan and Tsytovich.¹ As shown below, these disparities are evidently due to minor errors in their work.

In equation (3), β designates the species of the other particle with velocity \vec{v}_β , momentum \vec{p}_β , and background distribution $f_{p\beta}^{R(0)}$ involved in the bremsstrahlung process. Also, \vec{k} is the momentum transfer from the test particle and $E_k^{\sigma(0)}$ is the amplitude of the lowest order stochastic bremsstrahlung field with polarization \vec{e}_k^σ and mode σ . The matrix $\Lambda_{ij}^{(\beta)}(k, -\kappa)$ in equation (3) is defined by¹⁻³

$$\Lambda_{ij}^{(\alpha)}(k_1, k) = \frac{e_\alpha}{\gamma_\alpha m_\alpha} \left(\delta_{ij} + \frac{v_{\alpha i} k_j - v_{\alpha j} k_{1i}}{\omega - \vec{k} \cdot \vec{v}_\alpha - i\delta} - \frac{v_{\alpha i} v_{\alpha j} \left(\vec{k} \cdot \vec{k}_1 - \frac{\omega \omega_1}{c^2} \right)}{(\omega - \vec{k} \cdot \vec{v}_\alpha - i\delta)^2} \right). \quad (4)$$

The quantity $G_{mn}(k)$ in equation (3) is the linear photon Green's function, whose inverse is given by⁴

$$G_{ij}^{-1} = \frac{1}{\mu_0(\omega + i\delta)^2} (k_i k_j - k^2 \delta_{ij}) + \epsilon_{ij}. \quad (5)$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

²H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second-Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1995 (August 1983).

³H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1994 (September 1983).

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

For a spatially isotropic system, the dielectric permittivity tensor ϵ_{ij} is given by⁴⁻⁷

$$\epsilon_{ij}(\vec{k}, \omega) = \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) \epsilon_t(\vec{k}, \omega) + \frac{k_i k_j}{k^2} \epsilon_l(\vec{k}, \omega) , \quad (6)$$

where ϵ_t and ϵ_l are the transverse and longitudinal permittivity, respectively. In this case the linear photon Green's function is given by⁴⁻⁶

$$G_{ij} = \frac{k_i k_j}{k^2 \epsilon_l} + \frac{\omega^2}{\epsilon_t(\omega + i\delta)^2 - \frac{k^2}{\mu_0}} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) , \quad (7)$$

and the transverse and longitudinal permittivities are given by

$$\epsilon_t = \epsilon_0 + \frac{1}{2} \frac{1}{(\omega + i\delta)^2} \sum_s \frac{e_s^2}{k^2} \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \left[k^2 v_{sm} + \frac{k^2 v_s^2 - \omega \vec{k} \cdot \vec{v}_s}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} k_m \right] \frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} \quad (8)$$

and

$$\epsilon_l = \epsilon_0 + \frac{1}{\omega + i\delta} \sum_s \frac{e_s^2}{k^2} \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{1}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \vec{k} \cdot \vec{v}_s \vec{k} \cdot \vec{v}_{ps} f_{ps}^{R(0)} , \quad (9)$$

respectively. Here ϵ_0 and μ_0 are the permittivity and permeability, respectively, of the vacuum. Each species s of particles present in the system makes a contribution to equations (8) and (9). If the distribution function depends only on energy, then using the expression ϵ_s for the relativistic energy of a particle of species s , namely,

$$\epsilon_s = (p_s^2 c^2 + m_s^2 c^4)^{1/2} , \quad (10)$$

allows equations (8) and (9) to be easily rewritten in terms of $\partial f_{ps}^{R(0)} / \partial \epsilon_s$.^{4,5}

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

⁵A. I. Akhiezer, I. A. Akhiezer, R. V. Polovin, A. G. Sitenko, and K. N. Stepanov, *Plasma Electrodynamics, Vol. 1, Linear Theory*, Pergamon Press (1975), 206.

⁶V. N. Tsytovich, *Nonlinear Effects in Plasma*, Plenum Press, New York (1970), 314.

⁷V. N. Tsytovich, *Theory of Turbulent Plasma* (Consultants Bureau), Plenum Publishing Corp., New York (1977), 63-65.

In the present work, equations (2) and (3) are derived from first principles for the purpose of identifying some of the important physical assumptions and mathematical techniques involved in Tsytovich's theory of nonlinear bremsstrahlung and radiative instability in relativistic nonequilibrium beam-plasma systems (see also Selected Bibliography).¹ This will facilitate understanding and illustrate the procedures for reducing other components of the total nonlinear recoil force on a test particle due to bremsstrahlung emission.¹

The present work, together with other related work by the author (see also Selected Bibliography),²⁻⁴ is important for ongoing work in calculating collective radiative processes and conditions for the occurrence of radiative instability in relativistic nonequilibrium beam-plasma systems.

2. REDUCTION OF A COMPONENT OF THE BREMSSTRAHLUNG RECOIL FORCE

In this section, equation (3) is to be derived by reducing that part of equation (1) which corresponds to bremsstrahlung recoil. Also the quantity $\langle \vec{p}_\alpha^{\text{rad}(1b)} \rangle$ appearing in equation (2) will be defined. It has been shown by using the balance equations that the general form for the collective bremsstrahlung recoil force on a test particle participating in induced bremsstrahlung in a nonequilibrium beam-plasma system is given by^{1,8}

$$\vec{F}_\alpha^\sigma = -4\pi^3 \int \frac{d^3\vec{k} d\omega d^3\vec{k}' d^3\vec{p}_\beta}{(2\pi)^9} \frac{1}{\omega^2} \frac{\partial}{\partial \omega} (\omega^2 \epsilon^\sigma(\vec{k}, \omega)) \vec{k} (\vec{k} - \vec{k}') \cdot (\vec{v}_{p_\beta} f_{p_\beta}) \quad (11)$$

$$\times |E_k^{(0)}|^2 v_{p_\alpha, p_\beta}^\sigma(\vec{k}, \vec{k}') \delta(\omega - \vec{k} \cdot \vec{v}_\alpha + (\vec{k} - \vec{k}') \cdot \vec{v}_\beta) .$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

²H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second-Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1995 (August 1983).

³H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1994 (September 1983).

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

⁸H. E. Brandt, *On the Form of the Collective Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories HDL-TR-2026 (January 1984).

Here the dielectric constant $\epsilon^\sigma(\vec{k}, \omega)$ for the bremsstrahlung mode σ is defined in terms of the dielectric permittivity tensor $\epsilon_{ij}(\vec{k}, \omega)$ and the unit electric polarization vector \vec{e}_k^σ by^{4, 6, 8, *}

$$\epsilon^\sigma(\vec{k}, \omega) = e_{ki}^{\sigma*} \epsilon_{ij} e_{kj}^\sigma + \epsilon_0 \frac{c^2}{\omega^2} (\vec{k} \cdot \vec{e}_k^\sigma)(\vec{k} \cdot \vec{e}_k^{\sigma*}) . \quad (12)$$

Also in equation (11), $E_k^{\sigma(0)}$ is the zeroth-order amplitude of the stochastic bremsstrahlung field,⁴ and $V_{p\alpha, p\beta}^\sigma(\vec{k}, \vec{k})$ is the bremsstrahlung transition probability with the delta function factored out, expressing conservation of energy. It is clear that $V_{p\alpha, p\beta}^\sigma(\vec{k}, \vec{k})$ must be at least first order in the regular part of the field. In reducing equation (1), only those contributions which are of the same form as equation (11) are to be included, since only they correspond to bremsstrahlung.

The total field \vec{E}_k appearing in equation (1) and associated with the bremsstrahlung process involving the test particle is given by^{1, 4}

$$\vec{E}_k = \vec{E}_k^{\sigma(0)} + \vec{E}_k^{(1)} + \vec{E}_k^{(2)} + \vec{E}_{dpk}^{(1)} + \vec{E}_{dpk}^{(2)} . \quad (13)$$

Here $\vec{E}_k^{\sigma(0)}$ is the zeroth-order bremsstrahlung field

$$\vec{E}_k^{\sigma(0)} = E_k^{\sigma(0)} \vec{e}_k^\sigma , \quad (14)$$

where σ designates the mode and \vec{e}_k^σ is the polarization vector. If the bremsstrahlung field is to be real, then

$$\vec{E}_k^{\sigma(0)} = \vec{E}_{-k}^{\sigma(0)*} . \quad (15)$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

⁶V. N. Tsytovich, *Nonlinear Effects in Plasma*, Plenum Press, New York (1970), 314.

⁸H. E. Brandt, *On the Form of the Collective Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-2026 (January 1984)

*H. E. Brandt, *Bremsstrahlung Recoil Force on the Third-Order Nonlinear Dynamic Polarization Charge of a Relativistic Test Particle*, preprint, Harry Diamond Laboratories, 1983 (to be published).

The fields $\vec{E}_k^{(1)}$ and $\vec{E}_k^{(2)}$ in equation (13) are the self fields arising directly from the relativistic test particle's own motion, namely,⁴

$$\vec{E}_{kn}^{(1)} = \frac{-ie_\alpha}{(2\pi)^3(\omega + i\delta)} v_{\alpha m} G_{nm}(k) \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \quad (16)$$

and

$$\vec{E}_{kn}^{(2)} = \frac{-e_\alpha}{(2\pi)^3(\omega + i\delta)} G_{nm}(k) \quad (17)$$

$$\times \int \frac{dk_1}{\omega_1 - i\delta} E_{k_1 j}^* \Lambda_{mj}^{(\alpha)*}(k_1, k) \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_\alpha - \vec{k}_1 \cdot \vec{v}_\alpha) .$$

The fields $\vec{E}_{dpk}^{(1)}$ and $\vec{E}_{dpk}^{(2)}$ in equation (13) are the increasing order fields produced by the dynamic polarization current induced by the test particle and are given by⁴

$$\vec{E}_{dpkn}^{(1)} = \frac{i}{2(\omega + i\delta)} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2}{(\omega_1 - i\delta)(\omega_2 - i\delta)} \delta(k + k_1 + k_2) \quad (18)$$

$$\times [S_{mj}^{(s)}(k, -k_1, -k_2) + S_{mj}^{(s)}(k, -k_2, -k_1)] E_{k_1 j}^* E_{k_2 \ell}^*$$

and

$$\vec{E}_{dpkn}^{(2)} = \frac{-i}{\omega + i\delta} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \quad (19)$$

$$\times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)}(k, -k_1, -k_2, -k_3) E_{k_1 i}^* E_{k_2 j}^* E_{k_3 \ell}^* ,$$

respectively. The symmetrization in equation (19) appearing in earlier work^{1,4} has been removed. The delta functions appearing in equations (18) and (19) are four-dimensional, namely,

$$\delta(k) \equiv \delta^3(\vec{k}) \delta(\omega) . \quad (20)$$

The second-order nonlinear conductivity tensor $S_{ij\ell}^{(s)}(k, k_1, k_2)$ appearing in equation (18) is given by

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, *Harry Diamond Laboratories*, HDL-TR-1996 (September 1983).

$$S_{ijl}^{(s)}(k, k_1, k_2) = e_s^2 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \left[(\omega_1 - \vec{k}_1 \cdot \vec{v}_s) \frac{\partial}{\partial p_{sj}} + v_j k_{1m} \frac{\partial}{\partial p_{sm}} \right] \left[\frac{\partial}{\partial p_{sl}} + \frac{v_{sl}}{\omega_2 - \vec{k}_2 \cdot \vec{v}_s + i\delta} k_{2n} \frac{\partial}{\partial p_{sn}} \right] f_{ps}^{R(0)} . \quad (21)$$

The third-order nonlinear conductivity tensor $\Sigma_{ijlm}^{(s)}(k, k_1, k_2, k_3)$ is given by

$$\begin{aligned} \Sigma_{ijlm}^{(s)}(k, k_1, k_2, k_3) &= -ie_s^3 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \\ &\times [\delta_{jn}(\omega_1 - \vec{k}_1 \cdot \vec{v}_s) + k_{1n} v_{sj}] \frac{\partial}{\partial p_{sn}} \frac{1}{\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s + i\delta} \\ &\times [\delta_{lu}(\omega_2 - \vec{k}_2 \cdot \vec{v}_s) + k_{2u} v_{sl}] \frac{\partial}{\partial p_{su}} \frac{1}{\omega_3 - \vec{k}_3 \cdot \vec{v}_s + i\delta} \\ &\times [\delta_{mq}(\omega_3 - \vec{k}_3 \cdot \vec{v}_s) + k_{3q} v_{sm}] \frac{\partial}{\partial p_{sq}} f_{ps}^{R(0)} . \end{aligned} \quad (22)$$

Letting $B\vec{F}_\alpha^{(1)}$ denote the operation of including only that part of $\vec{F}_\alpha^{(1)}$ which is of the bremsstrahlung form given by equation (11), we obtain

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle = B \langle \vec{F}_\alpha^{(1)} \rangle . \quad (23)$$

Here $\langle \rangle$ denotes the ensemble average over the statistical phase distribution of the bremsstrahlung, which to the needed order is assumed to be random.⁴ Substituting equations (1) and (13) in equation (23) gives

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle = \sum_{n=1}^5 \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_n , \quad (24)$$

where

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_n = B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \cdot \vec{v}_\alpha \cdot \langle \vec{E}_k^n \rangle \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \quad (25)$$

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

and

$$\{\vec{E}_k^n, n = 1, 2, \dots, 5\} = \{\vec{E}_k^{\sigma(0)}, \vec{E}_k^{(1)}, \vec{E}_k^{(2)}, \vec{E}_{dpk}^{(1)}, \vec{E}_{dpk}^{(2)}\} . \quad (26)$$

To the needed order, the stochastic properties of the bremsstrahlung field are approximated by^{1,4}

$$\langle \vec{E}_{k1}^{\sigma(0)} \rangle = 0 \quad (27)$$

and

$$\langle \vec{E}_{k1}^{\sigma(0)} \vec{E}_{k1j}^{\sigma(0)} \rangle = e_{ki}^{\sigma} e_{kj}^{\sigma*} |\vec{E}_k^{\sigma(0)}|^2 \delta(k + k_1) . \quad (28)$$

Substituting equations (26) and (27) in equation (25) for $n = 1$ produces

$$\langle \vec{F}_\alpha^{\sigma grad(1)} \rangle_1 = 0 . \quad (29)$$

If we substitute equations (26) and (16) in equation (25) for $n = 2$, it is evident that the resulting expression is not of the required form given by equation (11). For example it does not contain $|\vec{E}_k^{\sigma(0)}|^2$; therefore the B operation gives zero, thus:

$$\langle \vec{F}_\alpha^{\sigma grad(1)} \rangle_2 = 0 . \quad (30)$$

Next substituting equations (26) and (17) in equation (25) for $n = 3$, we obtain

$$\begin{aligned} \langle \vec{F}_\alpha^{\sigma grad(1)} \rangle_3 = B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} v_{\alpha n} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) & \left[\frac{-e_\alpha G_{nm}(k)}{(2\pi)^3 (\omega + i\delta)} \right] \\ \times \int \frac{dk_1}{\omega_1 - i\delta} \langle \vec{E}_{k1j}^* \rangle \Lambda_{mj}^{(\alpha)*}(k_1, k) \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_\alpha - \vec{k}_1 \cdot \vec{v}_\alpha) . & \end{aligned} \quad (31)$$

¹A. V. Akopyan and V. N. Tsytoich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

Furthermore, substituting equation (13) in equation (31) and using equation (26) then

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_3 = \sum_{n=1}^5 \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{3n} , \quad (32)$$

where

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{3n} = & B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} v_{\alpha n} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \left[\frac{-e_\alpha G_{nm}(k)}{(2\pi)^3 (\omega + i\delta)} \right] \\ & \times \int \frac{dk_1}{\omega_1 - i\delta} \langle E_{k_1 j}^{n*} \rangle \Lambda_{mj}^{(\alpha)*}(k_1, k) \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_\alpha - \vec{k}_1 \cdot \vec{v}_\alpha) . \end{aligned} \quad (33)$$

Substituting equations (26) and (27) in equation (33) for $n = 1$ produces

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{31} = 0 . \quad (34)$$

Next substituting equations (26) and (16) in equation (33) for $n = 2$, we obtain an expression which is clearly not of the bremsstrahlung form given by equation (11), and therefore the B operation yields zero, thus:

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{32} = 0 . \quad (35)$$

Next substituting equations (26) and (17) in equation (33) for $n = 3$, we obtain a higher order Born term which is also not of the bremsstrahlung form given by equation (11), and therefore

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{33} = 0 . \quad (36)$$

For consistency with the Born approximation, only those terms which are first or second order in the regular part of the field are needed.¹

So that in the limit of $t \rightarrow \infty$ the integrals in equation (25) or equation (33) do not vanish, it is important to note here that the following combination of delta functions is needed in the integrand:

$$\delta^2(\omega - \vec{k} \cdot \vec{v}_\alpha) = \lim_{t \rightarrow \infty} \frac{t}{2\pi} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) . \quad (37)$$

In this way the t^{-1} factor is canceled. Equation (37) is obtained as follows. The well-known integral representation of the delta function is

$$\delta(\omega) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{-i\omega t} . \quad (38)$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

Therefore

$$\delta(\omega - \vec{k} \cdot \vec{v}) = \lim_{t \rightarrow \infty} \int_{-t/2}^{t/2} \frac{dt'}{2\pi} e^{-i(\omega - \vec{k} \cdot \vec{v})t'} \quad (39)$$

Then using equation (39) one has

$$\begin{aligned} \delta^2(\omega - \vec{k} \cdot \vec{v}_\alpha) &= \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \lim_{t \rightarrow \infty} \int_{-t/2}^{t/2} \frac{dt'}{2\pi} e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t'} \\ &= \lim_{t \rightarrow \infty} \int_{-t/2}^{t/2} \frac{dt'}{2\pi} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t'} \\ &= \lim_{t \rightarrow \infty} \int_{-t/2}^{t/2} \frac{dt'}{2\pi} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ &= \lim_{t \rightarrow \infty} \frac{t}{2\pi} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \end{aligned} \quad (40)$$

Thus in evaluating equation (31) one has an integrand containing the factor

$$F = \frac{1}{t} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \langle E_{k_1 j}^* \rangle \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_\alpha - \vec{k}_1 \cdot \vec{v}_\alpha) \quad (41)$$

or

$$F = \frac{1}{t} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \langle E_{k_1 j}^* \rangle \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \quad ; \quad (42)$$

or substituting equation (13) in equation (42) produces

$$F = \frac{1}{t} \sum_{n=1}^5 \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \langle E_{k_1 j}^{n*} \rangle \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \quad (43)$$

Because of equation (37), another factor of $\delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha)$ is needed to give a nonvanishing limit. For $n = 2$ in equation (43), one picks up a factor of $\delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha)$ by equation (16); however, the remaining integrand does not contain the factor $|E_k^{(0)}|^2$ necessary to be of the bremsstrahlung form given by equation (11). An iteration of equation (17) in equation (43) can give the quadratic delta function, but the remaining integrand is higher order. By analogous reasoning

$$\langle F_\alpha^{\text{grad}(1)} \rangle_{34} = 0 \quad , \quad (44)$$

$$\langle F_\alpha^{\text{grad}(1)} \rangle_{35} = 0 \quad . \quad (45)$$

Therefore, substituting equations (34) to (36), (44), and (45) in equation (32) produces

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_3 = 0 \quad (46)$$

The quantity $\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_4$ is nonvanishing and is calculated in a separate report.* Here the result of that calculation will be designated $\langle \vec{F}_\alpha^{\text{grad}(1b)} \rangle$. Thus

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_4 \equiv \langle \vec{F}_\alpha^{\text{grad}(1b)} \rangle \quad (47)$$

In the present report we emphasize the calculation of $\langle \vec{F}^{\text{grad}(1a)} \rangle$ in equation (2).

Substituting equations (26) and (19) in equation (25) one has

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_5 = & B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha n} \\ & \times \frac{-i}{\omega + i\delta} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \\ & \times \delta(k + k_1 + k_2 + k_3) \Gamma_{mij}^{(s)}(k, -k_1, -k_2, -k_3) \langle E_{k_1 i}^* E_{k_2 j}^* E_{k_3 l}^* \rangle \end{aligned} \quad (48)$$

One has, using equation (13) to the required order in equation (48),

$$\begin{aligned} \langle E_{k_1 i} E_{k_2 j} E_{k_3 l} \rangle = & \langle (E_{k_1 i}^{\sigma(0)} + E_{k_1 i}^{(1)} + E_{k_1 i}^{(2)} + E_{dpki}^{(1)} + E_{dpki}^{(2)}) \\ & \times (E_{k_2 j}^{\sigma(0)} + E_{k_2 j}^{(1)} + E_{k_2 j}^{(2)} + E_{dpkj}^{(1)} + E_{dpkj}^{(2)}) \\ & \times (E_{k_3 l}^{\sigma(0)} + E_{k_3 l}^{(1)} + E_{k_3 l}^{(2)} + E_{dpkl}^{(1)} + E_{dpkl}^{(2)}) \rangle \end{aligned} \quad (49)$$

If equation (49) is substituted in equation (48), the only possible nonvanishing contributions to the required order are given by

*H. E. Brandt, *Collective Bremsstrahlung Recoil Force on the Bare Charge of an Unperturbed Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, 1983 (to be published as HDL technical report).

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_5$$

$$\begin{aligned}
&= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha p} \\
&\times \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \\
&\times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)}(k, -k_1, -k_2, -k_3) \left\{ \frac{1}{2} E_{k3\ell}^{(1)*} \langle E_{k1i}^{\sigma(0)*} E_{k2j}^{\sigma(0)*} \rangle \right. \\
&\left. + \frac{1}{2} E_{k2j}^{(1)*} \langle E_{k1i}^{\sigma(0)*} E_{k3\ell}^{\sigma(0)*} \rangle + \frac{1}{2} E_{k1i}^{(1)*} \langle E_{k2j}^{\sigma(0)*} E_{k3\ell}^{\sigma(0)*} \rangle \right\} . \quad (50)
\end{aligned}$$

The factors of 1/2 in the three terms in the braces arise from the fact that equation (50) is effectively a self force; the field $E_{kn}^{(1)}$ in equation (50) and given by equation (16) is a field produced by the test particle α , and through the dynamic polarization that it induces it produces a force which acts back on the same test particle. Equation (50) may be rewritten as

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_5 = \sum_{n=1}^3 \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{5n} , \quad (51)$$

where

$$\begin{aligned}
&\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{51} \\
&= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha p} \\
&\times \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \\
&\times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)}(k, -k_1, -k_2, -k_3) \frac{1}{2} E_{k3\ell}^{(1)*} \langle E_{k1i}^{\sigma(0)*} E_{k2j}^{\sigma(0)*} \rangle \quad (52)
\end{aligned}$$

$$\langle \vec{F}_\alpha^{\sigma \text{rad}(1)} \rangle_{52}$$

$$= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha p} \\ \times \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \int_S e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \quad (53) \\ \times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)} \ell(k, -k_1, -k_2, -k_3) \frac{1}{2} E_{k_2j}^{(1)*} \langle E_{k_1i}^{\sigma(0)*} E_{k_3\ell}^{\sigma(0)*} \rangle$$

and

$$\langle \vec{F}_\alpha^{\sigma \text{rad}(1)} \rangle_{53}$$

$$= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha p} \\ \times \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \int_S e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \quad (54) \\ \times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)} \ell(k, -k_1, -k_2, -k_3) \frac{1}{2} E_{k_1i}^{(1)*} \langle E_{k_2j}^{\sigma(0)*} E_{k_3\ell}^{\sigma(0)*} \rangle$$

Substituting equations (28) and (16) in equation (52) produces

$$\langle \vec{F}_\alpha^{\sigma \text{rad}(1)} \rangle_{51}$$

$$= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ \times v_{\alpha p} \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \int_S e_s \int \frac{dk_1 dk_2 dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \quad (55) \\ \times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)} \ell(k, -k_1, -k_2, -k_3) \\ \times \frac{1}{2} \left[\frac{ie_\alpha}{(2\pi)^3 (\omega_3 - i\delta)} \right] v_{\alpha n} G_{ln}^*(k_3) \delta(\omega_3 - \vec{k}_3 \cdot \vec{v}_\alpha) e_{k_1i}^{\sigma*} e_{k_1j}^\sigma |E_{k_1}^{\sigma(0)}|^2 \\ \times \delta(k_1 + k_2)$$

Integrating equation (55) first over k_2 , using the property of the delta function $\delta(k_1 + k_2)$ and effectively setting $k_2 = -k_1$ (that is, $\vec{k}_2 = -\vec{k}_1$, $\omega_2 = -\omega_1$), we obtain

$$\begin{aligned}
\langle \vec{F}_\alpha^{\text{orad}(1)} \rangle_{51} &= B \lim_{t \rightarrow \infty} \frac{-e_\alpha^2}{8\pi^2 t} \int \frac{dk \, dk_1 \, dk_3}{(\omega + i\delta)^2 (\omega_1 - i\delta)(\omega_1 + i\delta)(\omega_3 - i\delta)^2} \vec{k} \delta(\vec{k} + \vec{k}_3) \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\
&\times v_{\alpha p} G_{pm}(k) \sum_S e_S \Sigma_{mlj}^{(S)}(k, -k_1, k_1, -k_3) \\
&\times v_{\alpha n} G_{ln}^*(k_3) \delta(\omega_3 - \vec{k}_3 \cdot \vec{v}_\alpha) e_{k_1 i}^{\sigma*} e_{k_1 j}^\sigma |E_{k_1}^{\sigma(0)}|^2 .
\end{aligned} \tag{56}$$

Next integrating equation (56) over k_3 produces

$$\begin{aligned}
\langle \vec{F}_\alpha^{\text{orad}(1)} \rangle_{51} &= B \lim_{t \rightarrow \infty} \frac{-e_\alpha^2}{8\pi^2 t} \int \frac{dk \, dk_1}{(\omega + i\delta)^4 \omega_1^2} \vec{k} \delta^2(\omega - \vec{k} \cdot \vec{v}_\alpha) \\
&\times v_{\alpha p} G_{pm}(k) \sum_S e_S \Sigma_{mlj}^{(S)}(k, -k_1, k_1, k) v_{\alpha n} G_{ln}^*(-k) \\
&\times e_{k_1 i}^{\sigma*} e_{k_1 j}^\sigma |E_{k_1}^{\sigma(0)}|^2 .
\end{aligned} \tag{57}$$

One also has

$$G_{nm}^*(-k) = G_{nm}(k) . \tag{58}$$

Equation (58) holds since⁴

$$E_{kn} = - \frac{i}{\omega + i\delta} G_{nm}(k) j_{km} \tag{59}$$

and by the reality of the current and field

$$E_{-kn}^* = E_{kn} \tag{60}$$

and

$$j_{-kn}^* = j_{kn} . \tag{61}$$

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

Taking the complex conjugate of equation (59), replacing k by $-k$, and substituting equations (60) and (61) produces

$$E_{kn} = \frac{-i}{\omega + i\delta} G_{nm}^*(-k) j_{km} \quad (62)$$

Comparing equations (59) and (62) shows that equation (58) follows. Next substituting equations (37) and (58) in equation (57), and noting that the t 's cancel, we find that the limit is trivial and

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{51} &= B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \int \frac{dk \, dk_1 \, \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha)}{(\omega + i\delta)^4 \omega_1^2} \\ &\times v_{\alpha p} G_{pm}(k) \sum_s e_s \Sigma_{mlj}^{(s)} \ell(k, -k_1, k_1, k) v_{\alpha n} G_{ln}(k) \\ &\times e_{k_1 i}^{\sigma*} e_{k_1 j}^\sigma |E_{k_1}^{\sigma(0)}|^2 \quad (63) \end{aligned}$$

Changing variables $\{k, k_1\}$ to $\{k_1, k\}$ and $\{i, j, \ell\}$ to $\{\ell, i, j\}$ in equation (63) produces

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{51} &= B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \sum_s e_s \int \frac{dk \, dk_1}{\omega^2 (\omega_1 + i\delta)^4} \vec{k}_1 |E_k^{\sigma(0)}|^2 \\ &\times v_{\alpha p} G_{pm}(k_1) v_{\alpha n} G_{jn}(k_1) e_{k\ell}^{\sigma*} e_{ki}^\sigma \\ &\times \Sigma_{mlj}^{(s)}(k_1, -k, k, k_1) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \quad (64) \end{aligned}$$

If we substitute equations (28) and (16) in equation (53), we obtain

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{52} &= B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ &\times v_{\alpha p} \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \sum_s e_s \int \frac{dk_1 \, dk_2 \, dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \\ &\times \delta(k + k_1 + k_2 + k_3) \Sigma_{mij}^{(s)} \ell(k, -k_1, -k_2, -k_3) \frac{1}{2} \left[\frac{ie_\alpha}{(2\pi)^3 (\omega_2 - i\delta)} \right] \\ &\times v_{\alpha n} G_{jn}^*(k_2) \delta(\omega_2 - \vec{k}_2 \cdot \vec{v}_\alpha) e_{k_1 i}^{\sigma*} e_{k_1 \ell}^\sigma |E_{k_1}^{\sigma(0)}|^2 \delta(k_1 + k_3) \quad (65) \end{aligned}$$

If we integrate over k_3 and k_2 , equation (65) becomes

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{52} = & B \lim_{t \rightarrow \infty} \left(\frac{-e_\alpha^2}{8\pi^2 t} \right) \int \frac{dk \, dk_1 \, \vec{k}}{(\omega + i\delta)^4 \omega_1^2} \\ & \times \delta^2(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\alpha p} G_{pm}(k) \sum_S e_S \Sigma_{mi}^{(S)} \ell(k, -k_1, k, k_1) \\ & \times v_{\alpha n} G_{jn}^*(-k) e_{k_1 i}^{\sigma*} e_{k_1 \ell}^\sigma |E_{k_1}^{\sigma(0)}|^2. \end{aligned} \quad (66)$$

Next substituting equations (37) and (58) in equation (66) produces

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{52} = & B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \int \frac{dk \, dk_1 \, \vec{k}}{(\omega + i\delta)^4 \omega_1^2} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ & \times v_{\alpha p} G_{pm}(k) \sum_S e_S \Sigma_{mi}^{(S)} \ell(k, -k_1, k, k_1) v_{\alpha n} G_{jn}(k) \\ & \times e_{k_1 i}^{\sigma*} e_{k_1 \ell}^\sigma |E_{k_1}^{\sigma(0)}|^2. \end{aligned} \quad (67)$$

Changing variables $\{k_1, k\}$ to $\{k, k_1\}$ and $\{i, \ell\}$ to $\{\ell, i\}$ in equation (67), we obtain

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{52} = & B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \int \frac{dk \, dk_1 \, \vec{k}_1}{\omega^2 (\omega_1 + i\delta)^4} |E_k^{\sigma(0)}|^2 \\ & \times v_{\alpha p} G_{pm}(k_1) v_{\alpha n} G_{jn}(k_1) e_{k \ell}^{\sigma*} e_{k i}^\sigma \\ & \times \sum_S e_S \Sigma_{mi}^{(S)} \ell(k_1, -k, k_1, k) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha). \end{aligned} \quad (68)$$

Next, substituting equations (28) and (16) in equation (54), we get

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{53} = & B \lim_{t \rightarrow \infty} \frac{2\pi}{t} e_\alpha \int \frac{dk}{\omega + i\delta} \vec{k} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ & \times v_{\alpha p} \left(\frac{-i}{\omega + i\delta} \right) G_{pm}(k) \sum_S e_S \int \frac{dk_1 \, dk_2 \, dk_3}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \\ & \times \delta(k + k_1 + k_2 + k_3) \Sigma_{mi}^{(S)} \ell(k, -k_1, -k_2, -k_3) \\ & \times \frac{1}{2} \left[\frac{ie_\alpha}{(2\pi)^3 (\omega_1 - i\delta)} \right] v_{\alpha n} G_{in}^*(k_1) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \\ & \times e_{k_2 j}^{\sigma*} e_{k_2 \ell}^\sigma |E_{k_2}^{\sigma(0)}|^2 \delta(k_2 + k_3). \end{aligned} \quad (69)$$

If we integrate over k_3 and k_1 , equation (69) becomes

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{53} &= B \lim_{t \rightarrow \infty} \left(\frac{-e_\alpha^2}{8\pi^2 t} \right) \int \frac{dk \, dk_2 \, \vec{k}}{(\omega + i\delta)^4 \omega_2^2} \\ &\times \delta^2(\omega - \vec{k} \cdot \vec{v}_\alpha) v_{\text{op}} G_{\text{pm}}(k) \sum_s e_s \Sigma_{\text{mj}}^{(s)} \ell(k, k, -k_2, k_2) \\ &\times v_{\text{om}} G_{\text{in}}^*(-k) e_{k_2 j}^{\sigma*} e_{k_2 \ell}^\sigma |E_{k_2}^{\sigma(0)}|^2. \end{aligned} \quad (70)$$

Next substituting equations (37) and (58) in equation (70) produces

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{53} &= B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \int \frac{dk \, dk_2 \, \vec{k}}{(\omega + i\delta)^4 \omega_2^2} \delta(\omega - \vec{k} \cdot \vec{v}_\alpha) \\ &\times v_{\text{op}} G_{\text{pm}}(k) \sum_s e_s \Sigma_{\text{mj}}^{(s)} \ell(k, k, -k_2, k_2) \\ &\times v_{\text{om}} G_{\text{in}}(k) e_{k_2 j}^{\sigma*} e_{k_2 \ell}^\sigma |E_{k_2}^{\sigma(0)}|^2. \end{aligned} \quad (71)$$

Changing variables $\{k, k_2\}$ to $\{k_1, k\}$ and $\{i, j, \ell\}$ to $\{j, \ell, i\}$ in equation (71) produces

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{53} &= B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \int \frac{dk \, dk_1 \, \vec{k}_1}{\omega^2 (\omega_1 + i\delta)^4} |E_k^{\sigma(0)}|^2 \\ &\times v_{\text{op}} G_{\text{pm}}(k_1) v_{\text{om}} G_{\text{jn}}(k_1) e_{k \ell}^{\sigma*} e_{ki}^\sigma \\ &\times \sum_s e_s \Sigma_{\text{mj}}^{(s)} \ell_i(k_1, k_1, -k, k) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha). \end{aligned} \quad (72)$$

However, equation (72) is not of the bremsstrahlung form given by equation (11); in particular, it does not contain the delta function factor $\delta(\omega - \vec{k} \cdot \vec{v}_\alpha + (\vec{k} - \vec{k}_1) \cdot \vec{v}_\beta)$ in the integrand. Such a factor could only appear through the third-order nonlinear conductivity tensor appearing in equation (72). According to equation (22), the factor $[\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s + i\delta]^{-1}$ appears in $\Sigma_{ij\ell m}^{(s)}(k, k_1, k_2, k_3)$, and it can be written

$$\begin{aligned} \frac{1}{\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s + i\delta} &= P \frac{1}{\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s} \\ &- i\pi \delta(\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s), \end{aligned} \quad (73)$$

and the imaginary part contains a delta function; however, in equation (72) it is $\Sigma_{mj\ell i}^{(s)}(k_1, k_1, -k, k)$ which appears, so that the delta function becomes $-i\pi\delta(0)$, which is not of the required form. Thus the B operation in equation (72) yields zero, and therefore

$$\langle \vec{F}_\alpha^{\text{grad}(1)} \rangle_{53} = 0 \quad (74)$$

Next substituting equations (64), (68), and (74) in equation (51) and the result together with equations (29), (30), (46), and (47) in equation (24), and comparing with equation (2), then

$$\begin{aligned} \langle \vec{F}_\alpha^{\text{grad}(1a)} \rangle = & B \left(\frac{-e_\alpha^2}{16\pi^3} \right) \sum_s e_s \int \frac{dk \, dk_1 \, \vec{k}_1}{\omega^2(\omega_1 + i\delta)^4} |E_k^{\sigma(0)}|^2 \\ & \times v_{\alpha p} G_{pm}(k_1) v_{\alpha n} G_{jn}(k_1) e_{k\ell}^* e_{ki}^\sigma \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \\ & \times [\Sigma_{m\ell ij}^{(s)}(k_1, -k, k, k_1) + \Sigma_{m\ell ji}^{(s)}(k_1, -k, k_1, k)] \quad (75) \end{aligned}$$

It remains to be shown in this report that equation (75) reduces to equation (3).

Evidently equation (75) should agree with equation (24) of Akopyan and Tsytovich.¹ Clearly there is a typographical error in the latter since a factor of e_α has been omitted. Also a factor of (-2) and complex conjugation signs on the conductivity tensor have apparently been omitted as shown below. Their equation (24)¹ contains an additional overall factor of $(4\pi)^2(2\pi)^{-6}$ due to the use there of Gaussian instead of MKS units and differing Fourier transform convention.

As was already noted elsewhere,⁴ extra factors of (4π) , $(2\pi)^{-6}$, and $(2\pi)^{-3}$ evidently enter through $E_{dpkn}^{(2)}$ in equation (19), due respectively to the use of Gaussian units, differing Fourier transform convention, and differing normalization of the background distribution $f_{ps}^{R(0)}$ in equation (21). However the normalization in equation (24) of Akopyan and Tsytovich¹ is apparently the same as that here. Otherwise an additional factor of $(2\pi)^{-3}$ was omitted there in equation (24). A factor of $(2\pi)^3$ arises if the normalization is restored. An additional factor of $(4\pi)(2\pi)^3$ enters through $E_{kn}^{(1)}$ in equation (16) due to the use of Gaussian units and differing Fourier transform convention. As already noted, an additional factor of $(2\pi)^{-3}$ enters through equation (1) due to the differing Fourier transform convention. These factors combine to

$$[(4\pi)(2\pi)^{-6}(2\pi)^{-3}][(2\pi)^3][4\pi(2\pi)^3][(2\pi)^{-3}] = (4\pi)^2(2\pi)^{-6} \quad .$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung In a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).

Because the same definition of $E_k^{\sigma(0)}$ in equation (28) here is used in equation (13) there, the differing Fourier transform convention is compensated for. Thus equation (75) is in essential agreement with equation (24) of Akopyan and Tsytovich¹ up to the omission of a factor of $(-2e_\alpha)$ and a complex conjugation sign on the conductivity tensor. This follows since the δ in $\omega_1 + i\delta$ may clearly be ignored, and

$$\Sigma_{mlij}^{(s)}(k_1, -k, k, k_1) = -iT_{mlij}^{(s)*}(k_1, k, -k, -k_1) \quad (76)$$

Also if the background is assumed to be nondissipative to the needed order then it follows that

$$(\omega_1 + i\delta)^{-1}G_{nj}(k_1) = [(\omega_1 + i\delta)^{-1}G_{jn}(k_1)]^* \quad (77)$$

which is true under certain conditions to be discussed below. Also for a spatially isotropic system the Green's function is symmetric in its indices according to equation (7). It should, however, be noted by comparing equation (14) of Akopyan and Tsytovich¹ with equation (21) of Brandt⁴ that the indices of G_{ij} are interchanged in Akopyan and Tsytovich¹ in the definition of the Green's function. It also follows that if one ignores the zero frequency resonance denominator, then equation (77) may be used provided the contribution of the imaginary part of the photonic Green's function is negligible to the needed order. Evidently the complex conjugation on the conductivity in equation (24) of Akopyan and Tsytovich¹ has been omitted. It is required if their equation (25) is to be consistent with the fact that the first and second complex denominators in their equation (21) are $(\omega - (\vec{k} \cdot \vec{v}_s) - i\delta)$ and $(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_s - i\delta)$, respectively.⁴ In equation (76) the tensor $T_{ijlm}^{(s)}(k, k_1, k_2, k_3)$ is given by^{1,4}

$$\begin{aligned} T_{ijlm}^{(s)}(k, k_1, k_2, k_3) = & e_s^3 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s - i\delta} \left[(\omega_1 - \vec{k}_1 \cdot \vec{v}_s) \frac{\partial}{\partial p_{sj}} \right. \\ & \left. + v_{sj}(\vec{k}_1 \cdot \vec{v}_{ps}) \right] \left\{ \frac{1}{\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_s - i\delta} \left[(\omega_2 - \vec{k}_2 \cdot \vec{v}_s) \frac{\partial}{\partial p_{sl}} \right. \right. \\ & \left. \left. + v_{sl}(\vec{k}_2 \cdot \vec{v}_{ps}) \right] \left[\frac{\partial}{\partial p_{sm}} + \frac{v_{sm}}{\omega_3 - \vec{k}_3 \cdot \vec{v}_s + i\delta} (\vec{k}_3 \cdot \vec{v}_{ps}) \right] \right\} f_{ps}^{R(0)} \quad (78) \end{aligned}$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung In a Nonequilibrium Relativistic Beam-Plasma System*, *Harry Diamond Laboratories*, HDL-TR-1996 (September 1983).

It was concluded elsewhere⁴ that for consistency, negative imaginary parts must be implicit in the first two denominators of equation (21) of Akopyan and Tsytovich.¹ By equation (22) here it is evident that

$$\Sigma_{ij}^{(S)} \ell_m(k_1, -k, k, k_1) = -\Sigma_{ij}^{(S)*} \ell_m(-k_1, k, -k, -k_1) . \quad (79)$$

Also from equations (22) and (78) it follows that

$$\Sigma_{ij}^{(S)} \ell_m(-k_1, k, -k, -k_1) = -iT_{ij}^{(S)} \ell_m(k_1, k, -k, -k_1) . \quad (80)$$

If we substitute equation (80) in equation (79), then equation (76) follows.

Equation (77) holds approximately if there is negligible linear absorption of energy by the background beam-plasma. To see this one notes that the energy $\epsilon_{abs}^{(0)}$ absorbed linearly by the background beam-plasma is given by

$$\epsilon_{abs}^{(0)} = \int d^3\vec{r} \int_{-\infty}^{\infty} dt \vec{E}(\vec{r}, t) \cdot \vec{j}(\vec{r}, t) , \quad (81)$$

where the Fourier transform of the field is given by equation (21) in an earlier report,⁴ namely

$$E_{ki} = \frac{-i}{\omega + i\delta} G_{ij}(k) j_{kj} . \quad (82)$$

In equation (82) G_{ij} is the linear photon Green's function of the background beam-plasma. Rewriting equation (81) in terms of the Fourier transformed quantities one obtains

$$\epsilon_{abs}^{(0)} = (2\pi)^4 \int dk E_{ki} \cdot j_{-ki} . \quad (83)$$

Next substituting equation (82) in equation (83) produces

$$\epsilon_{abs}^{(0)} = -i(2\pi)^4 \int \frac{dk}{\omega + i\delta} G_{ij}(k) j_{kj} j_{-ki} . \quad (84)$$

If the current \vec{j} is to be real, then

$$\vec{j}_{-k} = \vec{j}_k^* . \quad (85)$$

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

⁴H. E. Brandt, *The Total Field in Collective Bremsstrahlung In a Nonequilibrium Relativistic Beam-Plasma System*, *Harry Diamond Laboratories*, HDL-TR-1996 (September 1983).

Substituting equation (85) in equation (84) results in

$$\epsilon_{abs}^{(0)} = -i(2\pi)^4 \int dk \left(\frac{G_{ij}(k)}{\omega + i\delta} \right) j_{kj} j_{ki}^* \quad (86)$$

Also, clearly, since the energy must be real, then

$$\epsilon_{abs}^{(0)*} = \epsilon_{abs}^{(0)} \quad (87)$$

Then using equation (87), we find

$$\epsilon_{abs}^{(0)} = \frac{1}{2} \left(\epsilon_{abs}^{(0)} + \epsilon_{abs}^{(0)*} \right) \quad (88)$$

But using equation (86)

$$\epsilon_{abs}^{(0)*} = i(2\pi)^4 \int \frac{dk}{\omega - i\delta} G_{ij}^*(k) j_{kj}^* j_{ki} \quad (89)$$

Or, if we rename dummy indices $\{i, j\}$ as $\{j, i\}$, equation (89) becomes

$$\epsilon_{abs}^{(0)*} = -i(2\pi)^4 \int dk \left(\frac{-G_{ji}(k)}{\omega + i\delta} \right)^* j_{kj} j_{ki}^* \quad (90)$$

Then substituting equations (86) and (90) in equation (88), we get

$$\epsilon_{abs}^{(0)} = -\frac{i}{2} (2\pi)^4 \int dk \left[\left(\frac{G_{ij}(k)}{\omega + i\delta} \right) - \left(\frac{G_{ji}(k)}{\omega + i\delta} \right)^* \right] j_{ki} j_{kj}^* \quad (91)$$

If linear absorption by the background is to be vanishing for arbitrary current j_k , then equation (91) must be vanishing, and therefore

$$\frac{G_{ij}(k)}{\omega + i\delta} = \left(\frac{G_{ji}(k)}{\omega + i\delta} \right)^* \quad (92)$$

in which case equation (77) immediately follows. It is to be stressed that equation (77) is at best an approximation since the background will not be completely linearly nondissipative. In summary then, equation (75) is in essential agreement with equation (24) of Akopyan and Tsytovich¹ except for a typographical error of omission of a factor of $-2e_a$ in the latter; also Akopyan and Tsytovich there implicitly assume principal values with respect to single-wave particle resonance denominators as well as the approximate nondissipation relation, equation (77). However it is to be stressed that equation (75) is independent of these assumptions.

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

Proceeding with the further reduction of equation (75), one first defines

$$\begin{aligned} \bar{\Sigma}_{mlij}(k_1, -k, k, k_1) \equiv & B \int_S e_s [\Sigma_{mlij}^{(s)}(k_1, -k, k, k_1) \\ & + \Sigma_{mlji}^{(s)}(k_1, -k, k_1, k)] . \end{aligned} \quad (93)$$

Again the B operation selects only the part leading to the bremsstrahlung form given by equation (11). Using equation (93) in equation (75) produces

$$\begin{aligned} \langle F_{\alpha}^{\text{grad}(1a)} \rangle = & B \left(\frac{-e_{\alpha}^2}{16\pi^3} \right) \int \frac{dk \, dk_1 \, \vec{k}_1}{\omega^2 (\omega_1 + i\delta)^4} |E_k^{\sigma(0)}|^2 \\ & \times v_{\alpha p} G_{pm}(k_1) v_{\alpha n} G_{jn}(k_1) e_{k\ell}^{\sigma*} e_{ki}^{\sigma} \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_{\alpha}) \\ & \times \bar{\Sigma}_{mlij}(k_1, -k, k, k_1) . \end{aligned} \quad (94)$$

If we substitute equation (73) in equation (22), then equation (93) becomes

$$\begin{aligned} \bar{\Sigma}_{mlij}(k_1, -k, k, k_1) = & B(-i) \int_S e_s^4 \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{v_{sm}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_s + i\delta} \\ & \times [(-\omega + \vec{k} \cdot \vec{v}_s) \nabla_{ps\ell} + v_{s\ell} (-\vec{k} \cdot \vec{v}_{ps})] \left[P \frac{1}{\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_s} \right. \\ & \left. - i\pi \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_s) \right] \left\{ [(\omega - \vec{k} \cdot \vec{v}_s) \nabla_{psi} + v_{si} (\vec{k} \cdot \vec{v}_{ps})] \right. \\ & \times \left[\nabla_{psj} + \frac{v_{sj}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_s + i\delta} (\vec{k}_1 \cdot \vec{v}_{ps}) \right] \\ & + [(\omega_1 - \vec{k}_1 \cdot \vec{v}_s) \nabla_{psj} + v_{sj} (\vec{k}_1 \cdot \vec{v}_{ps})] \\ & \left. \times \left[\nabla_{psi} + \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \vec{k} \cdot \vec{v}_{ps} \right] \right\} f_{ps}^{R(0)} . \end{aligned} \quad (95)$$

If we compare equations (94) and (95) with equation (11), it is evident that only the part of equation (95) proportional to $\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta})$ can lead to the bremsstrahlung form given by equation (11), which is proportional to $\delta(\omega - \vec{k} \cdot \vec{v}_{\alpha} + (\vec{k} - \vec{k}) \cdot \vec{v}_{\beta})$. Note that because of the factor $\delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_{\alpha})$ in equation (94) one can set $\omega_1 = \vec{k}_1 \cdot \vec{v}_{\alpha}$ in $\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta})$ and also rename the integration variable \vec{k}_1 as $-\vec{k}$ to obtain $\delta(\omega - \vec{k} \cdot \vec{v}_{\alpha} + (\vec{k} - \vec{k}) \cdot \vec{v}_{\beta})$. Therefore the B operation in equation (95) selects, out of the sum over s, only the term $s = \beta$, and also only that part proportional to $\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta})$. Therefore equation (95) becomes

$$\bar{\Sigma}_{mlij} = B(-i)e_{\beta}^4 \int \frac{d^3 p_{\beta}}{(2\pi)^3} \frac{v_{\beta m}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta} + i\delta} [(-\omega + \vec{k} \cdot \vec{v}_{\beta}) \nabla_{p\beta l} + v_{\beta l} (-\vec{k} \cdot \vec{v}_{p\beta})] (-i\pi) D_{ij}(k, k_1, \vec{p}_{\beta}) f_{p\beta}^{R(0)} , \quad (96)$$

where the operator $D_{ij}(k, k_1, \vec{p}_{\beta})$ is defined by

$$D_{ij}(k, k_1, \vec{p}_{\beta}) \equiv \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) \left\{ [(\omega - \vec{k} \cdot \vec{v}_{\beta}) \nabla_{p\beta i} + v_{\beta i} (\vec{k} \cdot \vec{v}_{p\beta})] \left[\nabla_{p\beta j} + \frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta} + i\delta} \vec{k}_1 \cdot \vec{v}_{p\beta} \right] + [(\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta}) \nabla_{p\beta j} + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{p\beta})] \left[\nabla_{p\beta i} + \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_{\beta} + i\delta} \vec{k} \cdot \vec{v}_{p\beta} \right] \right\} . \quad (97)$$

Simplifying equations (96) and (97) produces

$$\bar{\Sigma}_{mlij} = B\pi e_{\beta}^4 \int \frac{d^3 p_{\beta}}{(2\pi)^3} \frac{v_{\beta m}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta} + i\delta} [(\omega - \vec{k} \cdot \vec{v}_{\beta}) \nabla_{p\beta l} + v_{\beta l} \vec{k} \cdot \vec{v}_{p\beta}] D_{ij} f_{p\beta}^{R(0)} \quad (98)$$

and

$$D_{ij} = \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) \left\{ (\omega - \vec{k} \cdot \vec{v}_{\beta}) \frac{\partial^2}{\partial p_{\beta i} \partial p_{\beta j}} + (\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta}) \frac{\partial^2}{\partial p_{\beta j} \partial p_{\beta i}} + (\omega - \vec{k} \cdot \vec{v}_{\beta}) \nabla_{p\beta i} \frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta} + i\delta} (\vec{k}_1 \cdot \vec{v}_{p\beta}) + v_{\beta i} (\vec{k} \cdot \vec{v}_{p\beta}) \nabla_{p\beta j} + v_{\beta i} (\vec{k} \cdot \vec{v}_{p\beta}) \frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta} + i\delta} (\vec{k}_1 \cdot \vec{v}_{p\beta}) + (\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta}) \nabla_{p\beta j} \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_{\beta} + i\delta} (\vec{k} \cdot \vec{v}_{p\beta}) + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{p\beta}) \nabla_{p\beta i} + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{p\beta}) \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_{\beta} + i\delta} (\vec{k} \cdot \vec{v}_{p\beta}) \right\} , \quad (99)$$

where the arguments of $D_{ij}(k, k_1, \vec{p}_{\beta})$ are suppressed. If we use the property of the delta function in equation (99) to replace $\omega_1 - \vec{k}_1 \cdot \vec{v}_{\beta}$ by $-\omega + \vec{k} \cdot \vec{v}_{\beta}$, the first two terms cancel and equation (99) becomes

$$\begin{aligned}
D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left[(\omega - \vec{k} \cdot \vec{v}_\beta) \nabla_{P\beta i} \frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \vec{k}_1 \cdot \vec{v}_{P\beta} \right. \\
& + v_{\beta i} (\vec{k} \cdot \vec{v}_{P\beta}) \nabla_{P\beta j} + v_{\beta i} (\vec{k} \cdot \vec{v}_{P\beta}) \frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \\
& - (\omega - \vec{k} \cdot \vec{v}_\beta) \nabla_{P\beta j} \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} (\vec{k} \cdot \vec{v}_{P\beta}) \\
& \left. + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \nabla_{P\beta i} + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} (\vec{k} \cdot \vec{v}_{P\beta}) \right] .
\end{aligned} \tag{100}$$

Using the properties of the derivative operator and using the delta function, one may rewrite equation (100) as

$$\begin{aligned}
D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ (\omega - \vec{k} \cdot \vec{v}_\beta) \left[\nabla_{P\beta i} \left(\frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k}_1 \cdot \vec{v}_{P\beta} \right. \\
& - v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \nabla_{P\beta i} + v_{\beta i} (\vec{k} \cdot \vec{v}_{P\beta}) \nabla_{P\beta j} + v_{\beta i} (\vec{k} \cdot \vec{v}_{P\beta}) \left(\frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \vec{k}_1 \cdot \vec{v}_{P\beta} \\
& - (\omega - \vec{k} \cdot \vec{v}_\beta) \left[\nabla_{P\beta j} \left(\frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k} \cdot \vec{v}_{P\beta} - v_{\beta i} (\vec{k} \cdot \vec{v}_{P\beta}) \nabla_{P\beta j} \\
& \left. + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \nabla_{P\beta i} + v_{\beta j} (\vec{k}_1 \cdot \vec{v}_{P\beta}) \frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} (\vec{k} \cdot \vec{v}_{P\beta}) \right\} .
\end{aligned} \tag{101}$$

In equation (101) the second and seventh terms cancel and the third and sixth terms cancel. Further simplifying equation (101) produces

$$\begin{aligned}
D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ (\omega - \vec{k} \cdot \vec{v}_\beta) \left[\nabla_{P\beta i} \left(\frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k}_1 \cdot \vec{v}_{P\beta} \right. \\
& - (\omega - \vec{k} \cdot \vec{v}_\beta) \left[\nabla_{P\beta j} \left(\frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k} \cdot \vec{v}_{P\beta} \\
& + v_{\beta i} \vec{k} \cdot \vec{v}_{P\beta} \left(\frac{v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \vec{k}_1 \cdot \vec{v}_{P\beta} \\
& \left. + v_{\beta j} \left[\vec{k}_1 \cdot \vec{v}_{P\beta} \left(\frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k} \cdot \vec{v}_{P\beta} - 2\pi i \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta) v_{\beta i} v_{\beta j} \vec{k} \cdot \vec{v}_{P\beta} \vec{k}_1 \cdot \vec{v}_{P\beta} \right\}
\end{aligned} \tag{102}$$

where equation (73) has been used in obtaining the last term. The last term may however be dropped since it does not lead to the bremsstrahlung form, equation (11), and therefore it will not contribute to equation (94). In particular, because of the factor $\delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta)$, the needed factor $\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta)$ becomes $\delta(\omega - \vec{k} \cdot \vec{v}_\beta)$. Also, if we replace $\omega - \vec{k} \cdot \vec{v}_\beta$ by $-\omega_1 + \vec{k}_1 \cdot \vec{v}_\beta$ in the first and $\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta$ by $-\omega + \vec{k} \cdot \vec{v}_\beta$ in the third term, equation (102) can be rewritten as

$$\begin{aligned}
D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ \left(-\nabla_{p\beta i} v_{\beta j} \right. \right. \\
& - (\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta) v_{\beta j} \left[\nabla_{p\beta i} \left(\frac{1}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k}_1 \cdot \vec{v}_{p\beta} \\
& - (\omega - \vec{k} \cdot \vec{v}_\beta) \left[\nabla_{p\beta j} \left(\frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k} \cdot \vec{v}_{p\beta} \\
& + v_{\beta i} \left(\left(\frac{1}{-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta} \right) (\vec{k} \cdot \vec{v}_{p\beta} v_{\beta j}) + v_{\beta j} \left[\vec{k} \cdot \vec{v}_{p\beta} \left(\frac{1}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k}_1 \cdot \vec{v}_{p\beta} \right. \\
& \left. \left. + v_{\beta j} \left[\vec{k}_1 \cdot \vec{v}_{p\beta} \left(\frac{v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \right] \vec{k} \cdot \vec{v}_{p\beta} \right\} . \quad (103)
\end{aligned}$$

Using the relativistic relation

$$v_{\beta i} = p_{\beta i} c^2 [p_\beta^2 c^2 + (m_\beta c^2)^2]^{-1/2} \quad (104)$$

produces

$$\nabla_{p\beta j} v_{\beta i} = \frac{1}{\gamma_{\beta m\beta}} \left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) , \quad (105)$$

where

$$\gamma_\beta = \left[1 + \left(\frac{p_\beta}{m_\beta c} \right)^2 \right]^{1/2} . \quad (106)$$

Performing the differentiations in equation (103) using equation (105) produces

$$\begin{aligned}
D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ \left[-\frac{1}{\gamma_{\beta m\beta}} \left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) \right. \right. \\
& - \left(\frac{k_{1\ell} v_{\beta j}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} \right) \frac{1}{\gamma_{\beta m\beta}} \left(\delta_{\ell i} - \frac{v_{\beta i} v_{\beta \ell}}{c^2} \right) + \left(\frac{v_{\beta i} k_{\ell}}{-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \frac{1}{\gamma_{\beta m\beta}} \\
& \times \left(\delta_{\ell j} - \frac{v_{\beta \ell} v_{\beta j}}{c^2} \right) + \frac{v_{\beta i} k_{1\ell} k_{m\ell} v_{\beta j}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta)^2 \gamma_{\beta m\beta}} \left(\delta_{\ell m} - \frac{v_{\beta \ell} v_{\beta m}}{c^2} \right) \left. \right] \vec{k}_1 \cdot \vec{v}_{p\beta} \\
& + \left[-\frac{1}{\gamma_{\beta m\beta}} \left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) - \frac{v_{\beta i} k_{\ell}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \frac{1}{\gamma_{\beta m\beta}} \left(\delta_{\ell j} - \frac{v_{\beta \ell} v_{\beta j}}{c^2} \right) \right. \\
& \left. + \frac{v_{\beta j} k_{1\ell}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \frac{1}{\gamma_{\beta m\beta}} \left(\delta_{i\ell} - \frac{v_{\beta \ell} v_{\beta i}}{c^2} \right) + \frac{v_{\beta j} v_{\beta i} k_{1\ell} k_{m\ell}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2 \gamma_{\beta m\beta}} \left(\delta_{m\ell} - \frac{v_{\beta m} v_{\beta \ell}}{c^2} \right) \right. \\
& \left. \left. \left. \frac{v_{\beta m} v_{\beta \ell}}{c^2} \right) \right] \vec{k} \cdot \vec{v}_{p\beta} \right\} . \quad (107)
\end{aligned}$$

Using the delta function to replace $\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta$ by $-\omega + \vec{k} \cdot \vec{v}_\beta$ in the denominator, we can rewrite equation (107) as

$$\begin{aligned}
 D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ \frac{1}{\gamma_{\beta m \beta}(-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta)} \left[\left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) (\omega - \vec{k} \cdot \vec{v}_\beta) \right. \right. \\
 & - v_{\beta j} \left(k_{1i} - \frac{\vec{k}_1 \cdot \vec{v}_\beta v_{\beta i}}{c^2} \right) + v_{\beta i} \left(k_j - \frac{\vec{k} \cdot \vec{v}_\beta}{c^2} v_{\beta j} \right) \\
 & + \frac{v_{\beta j} v_{\beta i}}{-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta} \left(\vec{k} \cdot \vec{k}_1 - \frac{(\vec{k} \cdot \vec{v}_\beta)(\vec{k}_1 \cdot \vec{v}_\beta)}{c^2} \right) \left. \right] \vec{k}_1 \cdot \vec{v}_{p\beta} \\
 & - \frac{1}{\gamma_{\beta m \beta}(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)} \left[\left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) (\omega - \vec{k} \cdot \vec{v}_\beta) + v_{\beta i} \left(k_j - \frac{\vec{k} \cdot \vec{v}_\beta}{c^2} v_{\beta j} \right) \right. \\
 & - v_{\beta j} \left(k_{1i} - \frac{\vec{k}_1 \cdot \vec{v}_\beta v_{\beta i}}{c^2} \right) - \frac{v_{\beta j} v_{\beta i}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \left(\vec{k} \cdot \vec{k}_1 - \frac{(\vec{k} \cdot \vec{v}_\beta)(\vec{k}_1 \cdot \vec{v}_\beta)}{c^2} \right) \left. \right] \vec{k} \cdot \vec{v}_{p\beta} \left. \right\} . \quad (108)
 \end{aligned}$$

When terms are combined, equation (108) becomes

$$\begin{aligned}
 D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \left\{ - \frac{1}{\gamma_{\beta m \beta}} \left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) (\vec{k}_1 + \vec{k}) \cdot \vec{v}_{p\beta} \right. \\
 & - \frac{v_{\beta j} k_{1i}}{\gamma_{\beta m \beta}(-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta)} \vec{k}_1 \cdot \vec{v}_{p\beta} - \frac{\vec{k} \cdot \vec{v}_\beta}{\gamma_{\beta m \beta} c^2} v_{\beta i} v_{\beta j} \left(\frac{\vec{k}_1}{-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta} \right. \\
 & + \left. \frac{\vec{k}}{-\omega + \vec{k} \cdot \vec{v}_\beta - i\delta} \right) \cdot \vec{v}_{p\beta} + \frac{v_{\beta i} k_j}{\gamma_{\beta m \beta}} \left(\frac{\vec{k}_1}{-\omega + \vec{k} \cdot \vec{v}_\beta + i\delta} + \frac{\vec{k}}{-\omega + \vec{k} \cdot \vec{v}_\beta - i\delta} \right) \cdot \vec{v}_{p\beta} \\
 & + \frac{v_{\beta i} v_{\beta j}}{\gamma_{\beta m \beta}} \left(\vec{k}_1 \cdot \vec{k} - \frac{\vec{k} \cdot \vec{v}_\beta \vec{k}_1 \cdot \vec{v}_\beta}{c^2} \right) \left(\frac{\vec{k}_1}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} + \frac{\vec{k}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2} \right) \cdot \vec{v}_{p\beta} \\
 & + \frac{v_{\beta j} k_{1i}}{\gamma_{\beta m \beta}(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)} \vec{k} \cdot \vec{v}_{p\beta} \\
 & - \left. \frac{v_{\beta i} v_{\beta j} \vec{k}_1 \cdot \vec{v}_\beta}{\gamma_{\beta m \beta} c^2} \left(\frac{\vec{k}_1}{\omega - \vec{k} \cdot \vec{v}_\beta - i\delta} + \frac{\vec{k}}{\omega - \vec{k} \cdot \vec{v}_\beta + i\delta} \right) \cdot \vec{v}_{p\beta} \right\} . \quad (109)
 \end{aligned}$$

Further simplifying equation (109) produces

$$\begin{aligned}
 D_{ij} = & \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \frac{1}{v_{\beta m \beta}} \left\{ - \left(\delta_{ij} - \frac{v_{\beta i} v_{\beta j}}{c^2} \right) (\omega - \vec{k} \cdot \vec{v}_\beta)^2 \right. \\
 & + \left[v_{\beta j} k_{1i} - v_{\beta i} k_{1j} + \frac{v_{\beta i} v_{\beta j}}{c^2} \vec{v}_\beta \cdot (\vec{k} - \vec{k}_1) \right] (\omega - \vec{k} \cdot \vec{v}_\beta) \\
 & + v_{\beta i} v_{\beta j} \left(\vec{k}_1 \cdot \vec{k} - \frac{\vec{k} \cdot \vec{v}_\beta \vec{k}_1 \cdot \vec{v}_\beta}{c^2} \right) \Bigg\} \\
 & \times \left[\frac{\vec{k}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2} + \frac{\vec{k}_1}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} \right] \cdot \vec{v}_{p\beta} .
 \end{aligned} \tag{110}$$

Equation (110) may be rewritten as

$$\begin{aligned}
 D_{ij} = & -\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \frac{1}{v_{\beta m \beta}} \left\{ (\omega - \vec{k} \cdot \vec{v}_\beta)^2 \delta_{ij} \right. \\
 & + (\omega - \vec{k} \cdot \vec{v}_\beta) (v_{\beta i} k_{1j} - v_{\beta j} k_{1i}) \\
 & - \frac{v_{\beta i} v_{\beta j}}{c^2} [(\omega - \vec{k} \cdot \vec{v}_\beta)^2 + \vec{v}_\beta \cdot (\vec{k} - \vec{k}_1) (\omega - \vec{k} \cdot \vec{v}_\beta) \\
 & + \vec{k} \cdot \vec{k}_1 c^2 - \vec{k} \cdot \vec{v}_\beta \vec{k}_1 \cdot \vec{v}_\beta] \Bigg\} \left[\frac{\vec{k}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2} + \frac{\vec{k}_1}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} \right] \cdot \vec{v}_{p\beta} .
 \end{aligned} \tag{111}$$

Further simplifying equation (111) produces

$$\begin{aligned}
 D_{ij} = & -\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \frac{1}{v_{\beta m \beta}} \left\{ (\omega - \vec{k} \cdot \vec{v}_\beta)^2 \delta_{ij} \right. \\
 & + (\omega - \vec{k} \cdot \vec{v}_\beta) (v_{\beta i} k_{1j} - v_{\beta j} k_{1i}) - \frac{v_{\beta i} v_{\beta j}}{c^2} [\vec{k} \cdot \vec{k}_1 c^2 \\
 & + \omega(\omega - \vec{k}_1 \cdot \vec{v}_\beta - \vec{k} \cdot \vec{v}_\beta)] \Bigg\} \left[\frac{\vec{k}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2} + \frac{\vec{k}_1}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} \right] \cdot \vec{v}_{p\beta} .
 \end{aligned} \tag{112}$$

Using the delta function to replace $(\omega - \vec{k}_1 \cdot \vec{v}_\beta - \vec{k} \cdot \vec{v}_\beta)$ by $(-\omega_1)$ in equation (112), we obtain

$$\begin{aligned}
 D_{ij} = & -\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \frac{1}{v_{\beta m \beta}} \left[(\omega - \vec{k} \cdot \vec{v}_\beta)^2 \delta_{ij} \right. \\
 & + (\omega - \vec{k} \cdot \vec{v}_\beta) (v_{\beta i} k_{1j} - v_{\beta j} k_{1i}) - v_{\beta i} v_{\beta j} \left(\vec{k} \cdot \vec{k}_1 - \frac{\omega \omega_1}{c^2} \right) \Bigg] \\
 & \times \left[\frac{\vec{k}}{(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)^2} + \frac{\vec{k}_1}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} \right] \cdot \vec{v}_{p\beta} .
 \end{aligned} \tag{113}$$

Substituting equation (4) in equation (113) produces

$$D_{ij} = -\frac{1}{e_\beta} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \times [\Lambda_{ij}^{(\beta)*}(\vec{k}_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(\vec{k}_1, k) \vec{k}_1] \cdot \vec{v}_{p\beta} . \quad (114)$$

Next substituting equation (114) in equation (98) results in

$$\bar{\Sigma}_{mlij} = -B\pi e_\beta^3 \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \frac{v_{\beta m}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta + i\delta} [(\omega - \vec{k} \cdot \vec{v}_\beta) \nabla_{p\beta l} + v_{\beta l} \vec{k} \cdot \vec{v}_{p\beta}] \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \times [\Lambda_{ij}^{(\beta)*}(\vec{k}_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(\vec{k}_1, k) \vec{k}_1] \cdot \vec{v}_{p\beta} f_{p\beta}^{R(0)} . \quad (115)$$

Integrating equation (115) by parts, then (since the background distribution $f_{p\beta}^{R(0)}$ is vanishing at infinity), one obtains

$$\bar{\Sigma}_{mlij} = B\pi e_\beta^3 \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \left\{ -\nabla_{p\beta l} \left[\frac{(\omega - \vec{k} \cdot \vec{v}_\beta) v_{\beta m}}{-\omega_1 + \vec{k}_1 \cdot \vec{v}_\beta - i\delta} \right] - \nabla_{p\beta n} \left[\frac{v_{\beta m} v_{\beta l} k_n}{-\omega_1 + \vec{k}_1 \cdot \vec{v}_\beta - i\delta} \right] \right\} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \times [\Lambda_{ij}^{(\beta)*}(\vec{k}_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(\vec{k}_1, k) \vec{k}_1] \cdot \vec{v}_{p\beta} f_{p\beta}^{R(0)} . \quad (116)$$

Performing the derivatives in equation (116) using equation (105), and then using the delta function to replace $\omega_1 - \vec{k}_1 \cdot \vec{v}_\beta$ by $-\omega + \vec{k} \cdot \vec{v}_\beta$ in the denominators, we obtain

$$\bar{\Sigma}_{mlij} = B\pi e_\beta^3 \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \left\{ -\frac{1}{\gamma_{\beta m} \beta} \left(\delta_{ml} - \frac{v_{\beta m} v_{\beta l}}{c^2} \right) + \frac{k_n}{\omega - \vec{k} \cdot \vec{v}_\beta - i\delta} \frac{1}{\gamma_{\beta m} \beta} \right. \\ \times \left(\delta_{nl} - \frac{v_{\beta n} v_{\beta l}}{c^2} \right) v_{\beta m} + \frac{1}{\omega - \vec{k} \cdot \vec{v}_\beta - i\delta} k_{1n} \frac{1}{\gamma_{\beta m} \beta} \left(\delta_{nl} - \frac{v_{\beta n} v_{\beta l}}{c^2} \right) v_{\beta m} \\ - \frac{1}{\omega - \vec{k} \cdot \vec{v}_\beta - i\delta} \frac{1}{\gamma_{\beta m} \beta} \left(\delta_{mn} - \frac{v_{\beta m} v_{\beta n}}{c^2} \right) v_{\beta l} k_n - \frac{v_{\beta m}}{\omega - \vec{k} \cdot \vec{v}_\beta - i\delta} \\ \times \frac{1}{\gamma_{\beta m} \beta} \left(\delta_{nl} - \frac{v_{\beta n} v_{\beta l}}{c^2} \right) k_n + \frac{v_{\beta m} v_{\beta l} k_n k_{1r}}{(\omega - \vec{k} \cdot \vec{v}_\beta - i\delta)^2} \frac{1}{\gamma_{\beta m} \beta} \left(\delta_{nr} - \frac{v_{\beta n} v_{\beta r}}{c^2} \right) \Big\} \\ \times \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) [\Lambda_{ij}^{(\beta)*}(\vec{k}_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(\vec{k}_1, k) \vec{k}_1] \cdot \vec{v}_{p\beta} f_{p\beta}^{R(0)} . \quad (117)$$

Combining terms in equation (117) produces

$$\begin{aligned} \bar{\Sigma}_{mlij} = & B\pi e_{\beta}^3 \int \frac{d^3 \vec{p}_{\beta}}{(2\pi)^3} \frac{1}{V_{\beta m \beta}} \left[-\delta_{m\ell} - \frac{k_m v_{\beta \ell} - k_1 \ell v_{\beta m}}{\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta} \right. \\ & \left. + \frac{v_{\beta m} v_{\beta \ell}}{c^2} \left(1 + \frac{(\vec{k} - \vec{k}_1) \cdot \vec{v}_{\beta}}{\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta} + \frac{\vec{k} \cdot \vec{k}_1 c^2}{(\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta)^2} - \frac{\vec{k} \cdot \vec{v}_{\beta} \vec{k}_1 \cdot \vec{v}_{\beta}}{(\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta)^2} \right) \right] \\ & \times \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) [\Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(k_1, k) \vec{k}_1] \cdot \vec{v}_{\beta} f_{P\beta}^{R(0)} . \end{aligned} \quad (118)$$

Further simplifying equation (118), we obtain

$$\begin{aligned} \bar{\Sigma}_{mlij} = & B\pi e_{\beta}^3 \int \frac{d^3 \vec{p}_{\beta}}{(2\pi)^3} \frac{1}{V_{\beta m \beta}} \left[-\delta_{m\ell} - \frac{k_m v_{\beta \ell} - k_1 \ell v_{\beta m}}{\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta} \right. \\ & \left. + \frac{v_{\beta m} v_{\beta \ell}}{c^2 (\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta)^2} [\omega(\omega - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) + \vec{k} \cdot \vec{k}_1 c^2] \right] \\ & \times \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) [\Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k} \\ & + \Lambda_{ij}^{(\beta)}(k_1, k) \vec{k}_1] \cdot \vec{v}_{\beta} f_{P\beta}^{R(0)} . \end{aligned} \quad (119)$$

If we use the delta function in equation (119) to replace $(\omega - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta})$ by $(-\omega_1)$, then after rearranging terms, we obtain

$$\begin{aligned} \bar{\Sigma}_{mlij} = & -B\pi e_{\beta}^3 \int \frac{d^3 \vec{p}_{\beta}}{(2\pi)^3} \frac{1}{V_{\beta m \beta}} \left[\delta_{\ell m} + \frac{v_{\beta \ell} k_m - v_{\beta m} k_1 \ell}{\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta} \right. \\ & \left. - \frac{v_{\beta \ell} v_{\beta m}}{(\omega - \vec{k} \cdot \vec{v}_{\beta} - i\delta)^2} \left(\vec{k} \cdot \vec{k}_1 - \frac{\omega \omega_1}{c^2} \right) \right] \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) \\ & \times [\Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(k_1, k) \vec{k}_1] \cdot \vec{v}_{\beta} f_{P\beta}^{R(0)} . \end{aligned} \quad (120)$$

Next substituting equation (4) in equation (120) produces

$$\begin{aligned} \bar{\Sigma}_{mlij} = & -B\pi e_{\beta}^2 \int \frac{d^3 \vec{p}_{\beta}}{(2\pi)^3} \Lambda_{\ell m}^{(\beta)}(k_1, k) \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_{\beta}) \\ & \times [\Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(k_1, k) \vec{k}_1] \cdot \vec{v}_{\beta} f_{P\beta}^{R(0)} . \end{aligned} \quad (121)$$

Equation (121) is to be compared with equation (26) of Akopyan and Tsytovich.¹ Since the B operation projects out $s = \beta$ as discussed above, equation (93) may be written

$$\bar{\Sigma}_{m\ell ij}(k_1, -k, k, k_1) = \text{Be}_\beta[\Sigma_{m\ell ij}^{(\beta)}(k_1, -k, k, k_1) + \Sigma_{m\ell ji}^{(\beta)}(k_1, -k, k_1, k)] \quad (122)$$

Comparing equations (121) and (122), we find

$$\begin{aligned} & B[\Sigma_{m\ell ij}^{(\beta)}(k_1, -k, k, k_1) + \Sigma_{m\ell ji}^{(\beta)}(k_1, -k, k_1, k)] \\ &= -B\pi e_\beta \int \frac{d^3 p_\beta}{(2\pi)^3} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \\ & \quad \times \Lambda_{\ell m}^{(\beta)}(k_1, k) [\Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k} + \Lambda_{ij}^{(\beta)}(k_1, k) \vec{k}_1] \cdot \vec{v}_\beta f_{p_\beta}^{R(0)} \quad (123) \end{aligned}$$

By equation (4) it follows that²

$$\delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \Lambda_{\ell m}^{(\beta)}(k_1, k) = \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \Lambda_{m\ell}^{(\beta)*}(k, k_1) \quad (124)$$

Substituting equations (76) and (124) in equation (123), dividing by i , and taking the complex conjugate, one obtains

$$\begin{aligned} & B[T_{m\ell ij}^{(\beta)}(k_1, k, -k, -k_1) + T_{m\ell ji}^{(\beta)}(k_1, k, -k_1, -k)] \\ &= B\pi e_\beta \int \frac{d^3 p_\beta}{(2\pi)^3} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \Lambda_{m\ell}^{(\beta)}(k, k_1) [\Lambda_{ij}^{(\beta)}(k_1, k) \vec{k} \\ & \quad + \Lambda_{ij}^{(\beta)*}(k_1, k) \vec{k}_1] \cdot \vec{v}_\beta f_{p_\beta}^{R(0)} \quad (125) \end{aligned}$$

If one ignores single-wave particle resonances, the $i\delta$ in equation (4) may be ignored and $\Lambda_{ij}^{(\beta)}(k_1, k)$ becomes effectively real and

$$P_1 \Lambda_{ij}^{(\beta)*}(k_1, k) = P_1 \Lambda_{ij}^{(\beta)}(k_1, k) \quad (126)$$

where P_1 denotes the operation of taking principal values with respect to Cerenkov resonance denominators of the form $(\omega - \vec{k} \cdot \vec{v}_\beta + i\delta)$. The principal part of $\Lambda_{ij}^{(\beta)}(k_1, k)$ is real. The nonprincipal part is proportional to $\delta(\omega - \vec{k} \cdot \vec{v}_\beta)$ or $\delta'(\omega - \vec{k} \cdot \vec{v}_\beta)$ according to equations (4) and (73). Therefore it will not yield the correct bremsstrahlung form and may be dropped. Operating on both sides of equation (125) with P_1 and using equation (126), we obtain

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

²H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second-Order in the Total Force in a Nonequilibrium Beam-Plasma System*, *Harry Diamond Laboratories, HDL-TR-1995* (August 1983).

$$\begin{aligned}
& P_1 B [T_{m\ell ij}^{(\beta)}(k_1, k, -k, -k_1) + T_{m\ell ji}^{(\beta)}(k_1, k, -k_1, -k)] \\
& = P_1 B i \pi e_\beta \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \\
& \quad \times \Lambda_{m\ell}^{(\beta)}(k, k_1) \Lambda_{ij}^{(\beta)}(k_1, k) (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta f_{p\beta}^{R(0)} .
\end{aligned} \tag{127}$$

Equation (127) agrees with equation (26) of Akopyan and Tsytovich¹ except for k appearing interchanged with k_1 in $\Lambda_{m\ell}^{(\beta)}$ and $\Lambda_{ij}^{(\beta)}$. This is apparently an error in the cited work.¹ The operation $P_1 B$ is implicit there and the i is absent since the imaginary part has been taken (the coefficient of i) and the right-hand side of equation (127) here is purely imaginary.

Next substituting equation (121) in equation (94) results in

$$\begin{aligned}
\langle F_\alpha^{\text{grad}(1a)} \rangle & = B \frac{e_\alpha^2 e_\beta^2}{16\pi^2} \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \frac{dk dk_1}{\omega^2 (\omega_1 + i\delta)^4} \\
& \quad \times \vec{k}_1 |E_k^{\sigma(0)}|^2 \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) \\
& \quad \times v_{\alpha p} G_{pm}(k_1) \Lambda_{\ell m}^{(\beta)}(k_1, k) e_{k\ell}^{\sigma*} [e_{ki}^{\sigma} \Lambda_{ij}^{(\beta)*}(k_1, k) G_{jn}(k_1) v_{\alpha n} \vec{k} \\
& \quad + e_{ki}^{\sigma} \Lambda_{ij}^{(\beta)}(k_1, k) G_{jn}(k_1) v_{\alpha n} \vec{k}_1] \cdot \vec{v}_\beta f_{p\beta}^{R(0)} .
\end{aligned} \tag{128}$$

Using equations (92) and (124) in equation (128) produces

$$\begin{aligned}
\langle F_\alpha^{\text{grad}(1a)} \rangle & = B \frac{e_\alpha^2 e_\beta^2}{16\pi^2} \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \frac{dk dk_1}{\omega^2 (\omega_1 + i\delta)^2 \omega_1^2} \vec{k}_1 |E_k^{\sigma(0)}|^2 \\
& \quad \times \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_\beta) \delta(\omega_1 - \vec{k}_1 \cdot \vec{v}_\alpha) [e_{k\ell}^{\sigma} \Lambda_{m\ell}^{(\beta)}(k, k_1) G_{mp}(k_1) v_{\alpha p}]^* \\
& \quad \times [e_{ki}^{\sigma} \Lambda_{ji}^{(\beta)}(k, k_1) G_{jn}(k_1) v_{\alpha n} \vec{k} + e_{ki}^{\sigma} \Lambda_{ji}^{(\beta)*}(k, k_1) G_{jn}(k_1) v_{\alpha n} \vec{k}_1] \cdot \vec{v}_\beta f_{p\beta}^{R(0)} .
\end{aligned} \tag{129}$$

Integrating over ω_1 , letting $\vec{k}_1 = -\vec{k}$, and denoting

$$\kappa = (\vec{k} \cdot \vec{v}_\alpha, \vec{k}) , \tag{130}$$

then we find that equation (129) becomes

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

$$\begin{aligned}
\langle \vec{F}_\alpha^{\text{grad}(1a)} \rangle = & -B \frac{e_\alpha^2 e_\beta^2}{16\pi^2} \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \frac{d\omega d^3 \vec{k} d^3 \vec{\kappa}}{\omega^2 (\vec{\kappa} \cdot \vec{v}_\alpha - i\delta)^2 (\vec{\kappa} \cdot \vec{v}_\alpha)^2} |E_k^{\sigma(0)}|^2 \\
& \times \delta(\omega - \vec{k} \cdot \vec{v}_\alpha + (\vec{k} - \vec{\kappa}) \cdot \vec{v}_\beta) [e_{k\ell}^\sigma \Lambda_{m\ell}^{(\beta)}(k, -\kappa) G_{mp}(-\kappa) v_{ap}]^* \\
& \times [e_{ki}^\sigma \Lambda_{ji}^{(\beta)}(k, -\kappa) G_{jn}(-\kappa) v_{an} \vec{k} \\
& - e_{ki}^\sigma \Lambda_{ji}^{(\beta)*}(k, -\kappa) G_{jn}(-\kappa) v_{an} \vec{k}] \cdot \vec{v}_\beta f_{p\beta}^{R(0)}.
\end{aligned} \tag{131}$$

If Cerenkov resonance is ignored, equation (126) applies. This is justified since only the principal part of $\Lambda_{ij}^{(\beta)}(k, -\kappa)$ leads to the bremsstrahlung form, and the principal part is real. Also, ignoring the $i\delta$ in $(\vec{k} \cdot \vec{v}_\alpha - i\delta)^{-1}$, dropping the unneeded B operation, and making the change of indices from $\{i, j, n, p\}$ to $\{j, i, p, n\}$, we find that equation (131) becomes

$$\begin{aligned}
\langle \vec{F}_\alpha^{\text{grad}(1a)} \rangle = & \frac{e_\alpha^2 e_\beta^2}{16\pi^2} \int \frac{d^3 \vec{p}_\beta}{(2\pi)^3} \frac{d\omega d^3 \vec{k} d^3 \vec{\kappa}}{\omega^2 (\vec{\kappa} \cdot \vec{v}_\alpha)^4} |E_k^{\sigma(0)}|^2 \\
& \times \delta(\omega - \vec{k} \cdot \vec{v}_\alpha + (\vec{k} - \vec{\kappa}) \cdot \vec{v}_\beta) \\
& \times |e_{kj}^\sigma \Lambda_{ij}^{(\beta)}(k, -\kappa) G_{ip}(-\kappa) v_{ap}|^2 (\vec{k} - \vec{\kappa}) \cdot \vec{v}_\beta f_{p\beta}^{R(0)}.
\end{aligned} \tag{132}$$

Equation (132) is in fact equation (3), as was to be shown. It is to be stressed that to obtain this expression it was necessary to assume that the background is approximately nonabsorptive and spatially isotropic. However, these assumptions are not involved in equation (128). As already stated in the introduction, equation (27) of Akopyan and Tsytovich¹ is apparently in error; namely, a factor of 2 is omitted there and the arguments of $\Lambda_{ij}^{(\beta)}(-\kappa, k)$ are interchanged. The additional factors of $(4\pi)^2 (2\pi)^{-6}$ there are apparently due to the use of Gaussian units and different Fourier transform conventions, as already discussed in equation (75). Also the Green's function used there is symmetric.

3. CONCLUSION

A number of important physical assumptions and mathematical techniques have been identified which are involved in Tsytovich's theory of nonlinear bremsstrahlung and radiative instability in relativistic nonequilibrium beam-plasma systems. In particular, the stochastic average of the lowest-order Born approximation for the nonlinear force on the bare charge of a relativistic test particle, equation (1), has been reduced to the form given by equations (2), (3), (47), (25), and (26). In the result of Akopyan and Tsytovich,¹ their equation (27) which corresponds to equation (3) here, a factor of 2 is evidently omitted and the arguments of $\Lambda_{ij}^{(\beta)}$ are incorrectly interchanged. Equation (3) has been shown to hold provided the background is approximately nonabsorptive and spatially isotropic. However, the form given by equation (128) holds without these assumptions. Also, an important expression, equation (123), was obtained involving the third-order nonlinear conductivity tensor. Using this expression it was shown that in equation (26) of Akopyan and Tsytovich¹ the arguments of $\Lambda_{ij}^{(\beta)}$ are again incorrectly interchanged.

The present work will facilitate better understanding of the methods necessary to reduce the other components of the total nonlinear force due to bremsstrahlung emission.¹ The latter is needed to determine the nonlinear bremsstrahlung probability and the conditions for the occurrence of a bremsstrahlung radiative instability in nonequilibrium relativistic beam-plasma systems.

¹A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].

LITERATURE CITED

1. A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung in a Nonequilibrium Plasma, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.* 1 (1975), 371].
2. H. E. Brandt, Nonlinear Force on an Unpolarized Relativistic Test Particle to Second-Order in the Total Field in a Nonequilibrium Beam-Plasma System, Harry Diamond Laboratories, HDL-TR-1995 (August 1983).
3. H. E. Brandt, Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System, Harry Diamond Laboratories, HDL-TR-1994 (September 1983).
4. H. E. Brandt, The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System, Harry Diamond Laboratories, HDL-TR-1996 (September 1983).
5. A. I. Akhiezer, I. A. Akhiezer, R. V. Polovin, A. G. Sitenko, K. N. Stepanov, *Plasma Electrodynamics*, Vol. 1, Linear Theory, Pergamon Press (1975), 206.
6. V. N. Tsytovich, *Nonlinear Effects in Plasma*, Plenum Press, New York (1970), 314.
7. V. N. Tsytovich, *Theory of Turbulent Plasma* (Consultants Bureau), Plenum Publishing Corp., New York (1977), 63-65.
8. H. E. Brandt, On the Form of the Collective Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System, Harry Diamond Laboratories, HDL-TR-2026 (January 1984).

SELECTED BIBLIOGRAPHY

Other Work of Akopyan and Tsytovich

V. N. Tsytovich, Bremsstrahlung of a Relativistic Plasma, Tr. Fiz. Inst. Akad. Nauk SSSR 66 (1973), 191 [Proc. (TRUDY) of P. N. Lebedev Physics Inst., 66 (1975), 199].

A. V. Akopyan and V. N. Tsytovich, Transition Bremsstrahlung of Relativistic Particles, Zh. Eksp. Teor. Fiz., 71 (1976), 166 [Sov. Phys. JETP, 44 (1976), 87].

A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung of Relativistic Electrons in a Plasma in a Strong Magnetic Field, Zh. Eksp. Teor. Fiz., 72 (1977), 1824 [Sov. Phys. JETP, 45 (1977), 957].

A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung of Relativistic Electrons in Plasmas, Proc. of XIIIth International Conference on Phenomena in Ionized Gases, Physical Society of German Democratic Republic (1977), 917.

A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung Instability of Relativistic Electrons in a Plasma, Astrofizika, 13 (1977), 717 [Astrophysics, 13 (1977), 423].

V. Tsytovich, Collective Effects in Bremsstrahlung of Fast Particles in Plasmas, Comments, Plasma Phys. Conf. Fusion, 4 (1978), 73.

Other Work of Brandt

H. E. Brandt, Symmetries of the Nonlinear Conductivity for a Relativistic Turbulent Plasma, Harry Diamond Laboratories, HDL-TR-1927 (March 1981).

H. E. Brandt, Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma, Phys. Fluids, 24 (1981), 1760.

H. E. Brandt, Comment on Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma, Phys. Fluids, 25 (1982), 1922.

H. E. Brandt, Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, in Plasma Astrophysics Course and Workshop, organized by the International School of Plasma Physics, 27 August to 7 September 1981, Varenna (Como), Italy (European Space Agency ESA SP-161, November 1981) (also to be published by Pergamon), p 361.

H. E. Brandt, On the Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, Harry Diamond Laboratories, HDL-TR-1970 (February 1982).

SELECTED BIBLIOGRAPHY (Cont'd)

Other Work of Brandt

H. E. Brandt, The Gluckstern-Hull Formula for Electron-Nucleus Bremsstrahlung, Harry Diamond Laboratories, HDL-TR-1884 (May 1980).

H. E. Brandt, Symmetry of the Complete Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, J. Math. Phys., 24 (1983), 1332, 2250.

H. E. Brandt, On the Form of the Collective Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System, Harry Diamond Laboratories, HDL-TR-2026 (January 1984).

Other Work of Brandt in Preparation

H. E. Brandt, Collective Bremsstrahlung Recoil Force on the Bare Charge of an Unperturbed Test Particle in a Nonequilibrium Beam-Plasma System, Harry Diamond Laboratories, 1983 (to be published as HDL technical report).

H. E. Brandt, Bremsstrahlung Recoil Force on the Third-Order Nonlinear Dynamic Polarization Charge of a Relativistic Test Particle, Harry Diamond Laboratories, 1982 (to be published as HDL technical report).

Other related work prepared in preprint form will be published later and is available from the author.

DISTRIBUTION

ADMINISTRATOR
DEFENSE TECHNICAL INFORMATION CENTER
ATTN DTIC-DDA (12 COPIES)
CAMERON STATION, BUILDING 5
ALEXANDRIA, VA 22314

COMMANDER
US ARMY MATERIEL DEVELOPMENT & READINESS
COMMAND
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333

DIRECTOR
DEFENSE ADVANCED RESEARCH PROJECTS
AGENCY
ATTN R. GULLICKSON
ATTN L. C. MARQUET
ATTN R. SEPUCHA
1400 WILSON BLVD
ARLINGTON, VA 22209

DIRECTOR
DEFENSE COMMUNICATIONS AGENCY
WASHINGTON, DC 20305

DIRECTOR
DEFENSE INTELLIGENCE AGENCY
ATTN D. SPOHN
ATTN DT-1, NUCLEAR & APPLIED
SCIENCES DIV
ATTN ELECTRONIC WARFARE BRANCH
WASHINGTON, DC 20301

UNDER SECRETARY OF DEFENSE
FOR RESEARCH & ENGINEERING
ATTN DEPUTY UNDER SECRETARY
(RESEARCH & ADVANCED TECH)
ATTN D. L. LAMBERSON
WASHINGTON, DC 20301

OFFICE OF THE UNDER SECRETARY OF
DEFENSE FOR RESEARCH & ENGINEERING
DIR ENERGY TECHNOLOGY OFFICE
THE PENTAGON
WASHINGTON, DC 20301

DIRECTOR
DEFENSE NUCLEAR AGENCY
ATTN DEPUTY DIRECTOR
SCIENTIFIC TECHNOLOGY
ATTN G. BAKER
ATTN RAEV, ELECTRONIC VULNERABILITY
ATTN J. Z. FARBER
ATTN G. K. SOPER
ATTN V. VAN LINT
WASHINGTON, DC 20305

DIRECTOR
NATIONAL SECURITY AGENCY
ATTN TECHNICAL LIBRARY
ATTN F. BEDARD
FT MEADE, MD 20755

ASSISTANT SECRETARY OF THE ARMY (RDA)
ATTN DEP FOR SCI & TECH
WASHINGTON, DC 20301

OFFICE, DEPUTY CHIEF OF STAFF
FOR OPERATIONS & PLANS
DEPT OF THE ARMY
ATTN DAMO-SSN, NUCLEAR DIV
WASHINGTON, DC 20310

OFFICE OF THE DEPUTY CHIEF OF STAFF
FOR RESEARCH, DEVELOPMENT,
& ACQUISITION
DEPARTMENT OF THE ARMY
ATTN DIRECTOR OF ARMY RESEARCH,
M. E. LASSER
ATTN DAMA-CSS-N, NUCLEAR TEAM
ATTN DAMA-ARZ-O, F. D. VERDERAME
WASHINGTON, DC 20310

COMMANDER
US ARMY ABERDEEN PROVING GROUND
ATTN STEAP-TL, TECH LIB
ABERDEEN PROVING GROUND, MD 21005

COMMANDER
BALLISTIC MISSILE DEFENSE
ADVANCED TECHNOLOGY CENTER
ATTN D. SCHENK
PO BOX 1500
HUNTSVILLE, AL 35807

DIRECTOR
US ARMY BALLISTIC RESEARCH LABORATORY
ATTN DRDAR-TSB-S (STINFO)
ATTN D. ECCLESHALL
ATTN C. HOLLANDSWORTH
ATTN R. KREMENS
ABERDEEN PROVING GROUND, MD 21005

US ARMY ELECTRONICS TECHNOLOGY &
DEVICES LABORATORY
ATTN DELET-DD
ATTN N. WILSON
FT MONMOUTH, NJ 07703

US ARMY ERADCOM
ATTN C. M. DESANTIS
FT MONMOUTH, NJ 07703

COMMANDER
US ARMY FOREIGN SCIENCE &
TECHNOLOGY CENTER
FEDERAL OFFICE BLDG

DISTRIBUTION (Cont'd)

US ARMY FOREIGN SCIENCE &
TECHNOLOGY CENTER (CONT'D)
ATTN DRXST-SD, SCIENCES DIV
ATTN T. CALDWELL
220 7TH STREET, NE
CHARLOTTESVILLE, VA 22901

DIRECTOR
US ARMY MATERIEL SYSTEMS
ANALYSIS ACTIVITY
ATTN DRXSY-MP
ABERDEEN PROVING GROUND, MD 21005

COMMANDER
US ARMY MATERIALS & MECHANICS
RESEARCH CENTER
ATTN DRXMR-H, BALLISTIC MISSILE
DEF MATLS PROG OFC
WATERTOWN, MA 02172

COMMANDER
US ARMY MISSILE COMMAND
ATTN DRDMI-TR, PHYSICAL SCIENCES DIR
REDSTONE ARSENAL, AL 35809

COMMANDER
US ARMY MISSILE & MUNITIONS
CENTER & SCHOOL
ATTN ATSK-CTD-F
REDSTONE ARSENAL, AL 35809

ARMY RESEARCH OFFICE (DURHAM)
PO BOX 12211
ATTN H. ROBL
ATTN R. LONTZ
ATTN B. D. GUENTHER
ATTN TECH LIBRARY
RESEARCH TRIANGLE PARK, NC 27709

COMMANDER
US ARMY RSCH & STD GP (EUR)
ATTN CHIEF, PHYSICS & MATH BRANCH
FPO NEW YORK 09510

NAVAL AIR SYSTEMS COMMAND
ATTN LCDR G. BATES, PMS-405
WASHINGTON, DC 20361

NAVAL INTELLIGENCE SUPPORT CENTER
ATTN M. KOONTZ
4301 SUITLAND RD
SUITLAND, MD 20390

NAVAL MATERIAL COMMAND
ATTN T. HORWATH
2211 JEFFERSON DAVIS HWY
WASHINGTON, DC 20301

OFFICE OF NAVAL RESEARCH
ATTN C. ROBERSON
ATTN W. J. CONDELL
800 N. QUINCY ST
ARLINGTON, VA 22217

SUPERINTENDENT
NAVAL POSTGRADUATE SCHOOL
ATTN LIBRARY, CODE 2124
MONTEREY, CA 93940

DIRECTOR
NAVAL RESEARCH LABORATORY
ATTN 2600, TECHNICAL INFO DIV
ATTN 5540, LASER PHYSICS
ATTN 6000, MATL & RADIATION SCI & TE
ATTN B. RIPIN
ATTN L. A. COSBY
ATTN E. E. KEMPE
ATTN J. T. SCHRIEMPF
ATTN R. F. WENZEL
ATTN R. HETTCHE
ATTN J. GOLDEN
ATTN V. L. GRANATSTEIN
ATTN M. A. REED
ATTN R. K. PARKER
ATTN P. A. SPRANGLE
ATTN S. GOLD
ATTN C. A. KAPETANAKOS
ATTN S. AHN
ATTN T. P. COFFEY
ATTN R. JACKSON
ATTN I. M. VITKOVITSKY
ATTN M. FRIEDMAN
ATTN J. PASOUR
ATTN G. COOPERSTEIN
WASHINGTON, DC 20375

COMMANDER
NAVAL SURFACE WEAPONS CENTER
ATTN V. PUGLIELLI
ATTN DX-21, LIBRARY DIV
DAHLGREN, VA 22448

COMMANDER
NAVAL SURFACE WEAPONS CENTER
ATTN J. Y. CHOE
ATTN F. SAZAMA
ATTN H. UHM
ATTN V. KENYON
ATTN E. NOLTING
ATTN WA-13, HIGH-ENERGY LASER BR
ATTN WA-50, NUCLEAR WEAPONS
EFFECTS DIV
ATTN WR, RESEARCH & TECHNOLOGY DEPT
ATTN WR-40, RADIATION DIV
ATTN E-43, TECHNICAL LIB
WHITE OAK, MD 20910

DISTRIBUTION (Cont'd)

COMMANDER
NAVAL WEAPONS CENTER
ATTN 315, LASER/INFRARED SYS DIV
ATTN 381, PHYSICS DIV
CHINA LAKE, CA 93555

HQ USAF/SAMI
WASHINGTON, DC 20330

ASSISTANT SECRETARY OF THE AIR FORCE
(RESEARCH & DEVELOPMENT)
WASHINGTON, DC 20330

DIRECTOR
AF OFFICE OF SCIENTIFIC RESEARCH
BOLLING AFB
ATTN NP, DIR OF PHYSICS
ATTN M. A. STROSCIO
ATTN R. BARKER
WASHINGTON, DC 20332

COMMANDER
AF WEAPONS LAB, AFSC
ATTN J. GENEROSA
ATTN A. H. GUENTHER
ATTN W. E. PAGE
ATTN LR, LASER DEV DIV
KIRTLAND AFB, NM 87117

WRIGHT-PATTERSON AFB
FOREIGN TECHNOLOGY DIVISION/ETD
ATTN J. BUTLER
WRIGHT-PATTERSON AFB, OH 45433

CENTRAL INTELLIGENCE AGENCY
ATTN R. PETTIS
ATTN D. B. NEWMAN
PO BOX 1925
WASHINGTON, DC 20013

DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
ATTN LIBRARY
ATTN CENTER FOR RADIATION RESEARCH
ATTN C. TEAGUE
ATTN E. MARX
WASHINGTON, DC 20234

DEPARTMENT OF ENERGY
ATTN O. P. MANLEY
WASHINGTON, DC 20585

NATIONAL AERONAUTICS & SPACE
ADMINISTRATION
ATTN R. RAMATY
GOODARD SPACE FLIGHT CENTER
GREENBELT, MD 20771

ADMINISTRATOR
NASA HEADQUARTERS
WASHINGTON, DC 20546

DIRECTOR
NASA
GODDARD SPACE FLIGHT CENTER
ATTN 250, TECH INFO DIV
GREENBELT, MD 20771

AMES LABORATORY (ERDA)
IOWA STATE UNIVERSITY
ATTN NUCLEAR SCIENCE CATEGORY
AMES, IA 50011

ASTRONOMICAL INSTITUTE--ZONNENBURG 2
ATTN M. KUPERUS
3512 NL UTRECHT, THE NETHERLANDS

THE AUSTRALIAN NATIONAL UNIVERSITY
DEPT OF THEORETICAL PHYSICS
ATTN D. B. MELROSE
PO BOX 4
CANBERRA A.C.T. 2600
AUSTRALIA

B. K. DYNAMICS
ATTN R. J. LINZ
15825 SHADY GROVE RD
ROCKVILLE, MD 20850

BOEING AEROSPACE
ATTN H. T. BUSHNER
PO BOX 3999 MS 8R-10
SEATTLE, WA 98124

BROOKHAVEN NATIONAL LABORATORY
ASSOCIATED UNIVERSITIES, INC
ATTN PHYSICS DEPT
UPTON, LONG ISLAND, NY 11973

CALIFORNIA INSTITUTE OF TECHNOLOGY
THEORETICAL ASTROPHYSICS
ATTN K. THORNE
130-33
PASADENA, CA 91125

UNIVERSITY OF CALIFORNIA
DEPT OF APPL PHYS & INFORMATION
SCIENCE
ATTN H. ALFVEN
LA JOLLA, CA 92093

CALIFORNIA INSTITUTE OF TECHNOLOGY
JET PROPULSION LABORATORY
ATTN J. C. HUBBARD
IRVINE, CA 92717

DISTRIBUTION (Cont'd)

UNIVERSITY OF CALIFORNIA
ELECTRICAL ENGINEERING DEPARTMENT
ATTN C. K. BIRDSALL
BERKELEY, CA 94720

UNIVERSITY OF CALIFORNIA
PHYSICS DEPT
ATTN A. N. KAUFMAN
BERKELEY, CA 94720

UNIVERSITY OF CALIFORNIA, DAVIS
APPLIED SCIENCES
ATTN J. DEGROOT
DAVIS, CA 95616

UNIVERSITY OF CALIFORNIA, IRVINE
DEPT OF PHYSICS
ATTN G. BENFORD
ATTN N. ROSTOKER
ATTN M. MAYER
IRVINE, CA 92717

UNIVERSITY OF CALIFORNIA, LOS ANGELES
DEPT OF PHYSICS
ATTN K. NOZAKI
LOS ANGELES, CA 90025

CAMBRIDGE UNIVERSITY
INSTITUTE OF ASTRONOMY
MADINGLEY ROAD
ATTN M. REES
CAMBRIDGE CB 3 0 HA, ENGLAND

CHALMERS UNIV OF TECHNOLOGY
INST OF ELECTROMAGNETIC FIELD
THEORY
ATTN H. WILHELMSSON
S-41296 GOTHENBURG, SWEDEN

UNIVERSITY OF CHICAGO
LAB FOR ASTROPHYSICS & SPACE RESEARCH
ATTN E. PARKER
CHICAGO, IL 60637

UNIVERSITY OF COLORADO
DEPT OF ASTROGEOPHYSICS
ATTN M. GOLDMAN
ATTN D. SMITH
BOULDER, CO 80309

COLUMBIA UNIVERSITY
ATTN S. JOHNSTON
216 MUDD BLDG
NEW YORK, NY 10027

CORNELL UNIVERSITY
ATTN R. LOVELACE
ATTN R. N. SUDAN
ATTN J. NATION
ATTN D. HAMMER

CORNELL UNIVERSITY (CONT'D)
ATTN H. FLEISHMANN
ITHACA, NY 14853

DARTMOUTH COLLEGE
PHYSICS DEPT
ATTN J. E. WALSH
HANOVER, NH 03755

UNIVERSITY OF THE DISTRICT
OF COLUMBIA
VAN NESS CAMPUS
DEPT OF PHYSICS
ATTN M. J. SMITH
4200 CONNECTICUT AVE, NW
WASHINGTON, DC 20008

ENGINEERING SOCIETIES LIBRARY
345 EAST 47TH STREET
ATTN ACQUISITIONS DEPARTMENT
NEW YORK, NY 10017

GENERAL DYNAMICS
ATTN K. H. BROWN
PO BOX 2507 MZ 44-21
POMONA, CA 91769

UNIVERSITY OF ILLINOIS AT
URBANA--CHAMPAIGN
DEPT OF PHYSICS
ATTN N. IWAMOTO
URBANA, IL 61801

IRT CORP
ATTN F. CHILTON
PO BOX 85317
SAN DIEGO, CA 92138

ISTITUTO DI FISICA DELL' UNIVERSITA
VIA CELORIA 16
ATTN P. CALDIROLA
ATTN C. PAIZIS
ATTN E. SINDONI
20133 MILANO, ITALY

ISTITUTO DI FISICA DELL' UNIVERSITA
ATTN A. CAVALIERE
ATTN R. RUFFINI
ROME, ITALY

ISTITUTO DI FISICA GENERALE
DELL' UNIVERSITA
CORSO M. D'AZEGLIO
ATTN A. FERRARI
46 TORINO, ITALY

JAYCOR
ATTN E. CONRAD
205 S. WHITING ST
ALEXANDRIA, VA 22304

DISTRIBUTION (Cont'd)

KAMAN SCIENCES CORP
ATTN T. A. TUMOLILLO
1500 GARDEN OF THE GODS
COLORADO SPRINGS, CO 80907

LAWRENCE BERKELEY LAB
ATTN A. FALTENS
ATTN A. M. SESSLER
ATTN D. KEEFE
BERKELEY, CA 94720

LAWRENCE LIVERMORE LABORATORY
UNIVERSITY OF CALIFORNIA
ATTN L. MARTIN
ATTN P. WHEELER
ATTN H. S. CABAYAN
ATTN R. BRIGGS
ATTN E. K. MILLER
ATTN R. ZIOLKOWSKI
ATTN R. SCARPETTI
ATTN R. MINICH
ATTN S. BURKHART
ATTN G. VOGTLIN
ATTN V. W. SLIVINSKY
ATTN S. L. YU
ATTN G. CRAIG
ATTN R. ALVAREZ
ATTN G. LASCHE
ATTN J. H. YEE
ATTN H. W. MELDNER
ATTN J. S. PETTIBONE
ATTN A. J. POGGIO
ATTN M. LOYD
PO BOX 808
LIVERMORE, CA 94550

CHIEF
LIVERMORE DIVISION, FIELD COMMAND, DNA
LAWRENCE LIVERMORE LABORATORY
PO BOX 808
LIVERMORE, CA 94550

LOS ALAMOS SCI LAB
ATTN S. COLGATE
ATTN D. F. DUBOIS
ATTN D. FORSLAND
ATTN C. M. FOWLER
ATTN B. WARNER
ATTN J. LANDT
ATTN R. HOEBERLING
ATTN T. R. KING
ATTN A. KADISH
ATTN K. LEE
ATTN R. W. FREYMAN
ATTN L. M. DUNCAN
ATTN D. B. HENDERSON
ATTN L. E. THODE
ATTN H. A. DAVIS
ATTN T. KWAN
PO BOX 1663
LOS ALAMOS, NM 87545

UNIVERSITY OF MARYLAND
DEPT OF ELECTRICAL ENGINEERING
ATTN M. REISER
ATTN W. DESTLER
ATTN C. D. STRIFFLER
ATTN M. T. RHEE
COLLEGE PARK, MD 20742

UNIVERSITY OF MARYLAND
DEPT OF PHYSICS & ASTRONOMY
ATTN H. R. GRIEM
ATTN E. OTT
ATTN C. GREBOGI
ATTN K. PAPADOPOULOS
ATTN J. WEBER
COLLEGE PARK, MD 20742

MASSACHUSETTS INSTITUTE
OF TECHNOLOGY
PLASMA FUSION CENTER
ATTN G. BEKEFI
ATTN J. BELCHER
ATTN T. CHANG
ATTN B. COPPI
ATTN R. DAVIDSON
ATTN S. OLBERT
ATTN B. ROSSI
CAMBRIDGE, MA 02139

MAX PLANCK INST, FUR AERONOMIE
3411 KATLENBG
ATTN J. AXFORD
ATTN E. MARSCH
LINDAU 3, WEST GERMANY

MAX PLANCK INSTITUTE OF PHYSICS
& ASTROPHYSICS
ATTN M. NORMAN
ATTN J. TRUMPER
GARCHING BEI MUNICH, WEST GERMANY

MAXWELL LABORATORIES
ATTN J. S. PEARLMAN
8835 BALBOA AVE
SAN DIEGO, CA 92123

MISSION RESEARCH CORP
ATTN B. GODFREY
ATTN D. J. SULLIVAN
ATTN M. BOLLEN
ATTN C. LONGMIRE
ATTN D. VOSS
1720 RANDOLPH RD, SE
ALBUQUERQUE, NM 87106

UNIVERSITY OF NAGOYA
DEPT OF PHYSICS
ATTN S. HAYAKAWA
NAGOYA, JAPAN

DISTRIBUTION (Cont'd)

NATIONAL RESEARCH COUNCIL
DIVISION OF PHYSICS
ATTN P. JAANIMAGI
OTTOWA, ONTARIO
CANADA

NATIONAL SCIENCE FOUNDATION
DIVISION OF ASTRONOMICAL SCIENCES
ATTN M. AIZENMAN
1800 G STREET, NW
WASHINGTON, DC 20550

OAK RIDGE NATIONAL LABORATORY
PO BOX Y
ATTN A. C. ENGLAND
OAK RIDGE, TN 37830

OCCIDENTAL RESEARCH CORP
ATTN D. B. CHANG
2100 SE MAIN
IRVINE, CA 92713

OXFORD UNIVERSITY
DEPARTMENT THEORETICAL PHYSICS
1 KEBLE RD
ATTN D. TER HAAR
OXFORD OX1 3NP, ENGLAND

OXFORD UNIVERSITY
ASTROPHYSICS DEPARTMENT
SOUTH PARK ROAD
ATTN A. HALL
OXFORD--OXI 3 RQ, ENGLAND

OXFORD UNIVERSITY
DEPT OF ENGINEERING SCIENCE
PARKS ROAD
ATTN L. M. WICKENS
ATTN J. E. ALLEN
OXFORD, UNITED KINGDOM

PHYSICS INTERNATIONAL
ATTN B. A. LIPPMANN
ATTN R. D. GENUARIO
2700 MERCER ST
SAN LEANDRO, CA 94577

PRINCETON UNIVERSITY
ASTROPHYSICAL SCIENCES
PETON HALL
ATTN R. KULSRUD
PRINCETON, NJ 08540

PRINCETON UNIVERSITY
PLASMAPHYSICS LABORATORY
ATTN R. WHITE
PRINCETON, NJ 08540

QUEEN MARY COLLEGE
DEPT OF APPLIED MATH
MILE END ROAD
ATTN D. BURGESS
LONDON E1 4NS, ENGLAND

RUHR UNIVERSITY
ATTN W. SEBOLD
ATTN W. ZWINGMANN
4630 BOCHUM 1, WEST GERMANY

INSTITUTE OF EXPERIMENTAL PHYSICS
RUHR-UNIVERSITY
POSTFACH 2148
ATTN H. KUNZE
436 BOCHUM, WEST GERMANY

SACHS/FREEMAN ASSOC
ATTN R. A. MAHAFFEY
19300 GALLANT FOX
SUITE 214
BOWIE, MD 20715

SANDIA LABORATORIES
ATTN R. B. MILLER
ATTN J. POWELL
ATTN W. BALLARD
ATTN C. EKDAHL
ATTN W. D. BROWN
ATTN S. HUMPHRIES, JR.
ATTN G. ROHWEIN
ATTN K. PRESTWICH
4244 SANDIA LABS
ALBUQUERQUE, NM 87115

SCIENCE APPLICATIONS, INC
ATTN E. P. CORNET
ATTN A. DROBOT
ATTN E. KANE
1710 GOODRIDGE DR
PO BOX 1303
MCLEAN, VA 22012

SCIENCE APPLICATIONS, INC
ATTN F. CHILTON
1200 PROSPECT ST
PO BOX 2351
LA JOLLA, CA 92038

SERVICE DE CHEMIE PHYSIQUELL
ATTN R. BALESCU
CAMPUS PLAINE U.L.B.
CODE POSTAL n° 231
BOULEVARD DU TRIOMPHE
10 50 BRUXELLES, BELGIUM

SPECOLA VATICANA
ATTN W. STOEGER, S. J.
I-00120 CITTA DEL VATICANO
ITALY

DISTRIBUTION (Cont'd)

SRI INTERNATIONAL
ATTN G. AUGUST
ATTN C. L. RINO
333 RAVENSWOOD AVE
MENLO PARK, CA 94025

STANFORD UNIVERSITY
INST PLASMA RES
ATTN P. A. STURROCK
STANFORD, CA 94305

STERRERWACHT-LEIDEN
ATTN C. A. NORMAN
ATTN P. ALLAN
2300 RA LEIDEN, THE NETHERLANDS

TEL-AVIV UNIVERSITY
DEPT OF PHYSICS & ASTRONOMY
ATTN G. TAUBER
TEL AVIV, ISRAEL

TELEDYNE BROWN ENGINEERING
CUMMINGS RESEARCH PARK
ATTN MELVIN L. PRICE, MS-44
HUNTSVILLE, AL 35807

INSTITUTE FOR THEORETICAL MECHANICS-
RUG
KRIJGSLAAN 271-S9
ATTN F. VERHEEST
B-9000 GENT, BELGIUM

UNIVERSITY OF TENNESSEE
DEPT OF ELECTRICAL ENGINEERING
ATTN I. ALEXEFF
ATTN J. R. ROTH
KNOXVILLE, TN 37916

UMEA UNIVERSITY
DEPT OF PLASMA PHYSICS
ATTN J. LARSSON
ATTN L. STENFLO
S-90187 UMEA
SWEDEN

UNIVERSITY OF WASHINGTON
DEPT OF PHYSICS
ATTN M. BAKER
SEATTLE, WA 98195

WEIZMANN INSTITUTE
DEPT OF NUCLEAR PHYSICS
ATTN AMRI WANDEL
REHOVOT, ISRAEL

WESTERN RESEARCH CORP
ATTN R. O. HUNTER
225 BROADWAY, SUITE 1600
SAN DIEGO, CA 92101

US ARMY ELECTRONICS RESEARCH
& DEVELOPMENT COMMAND
ATTN COMMANDER, DRDEL-CG
ATTN TECHNICAL DIRECTOR, DRDEL-CT
ATTN PUBLIC AFFAIRS OFFICE, DRDEL-IN

HARRY DIAMOND LABORATORIES
ATTN D/TSO/DIVISION DIRECTORS
ATTN RECORD COPY, 81200
ATTN HDL LIBRARY, 81100 (3 COPIES)
ATTN HDL LIBRARY (WOODBIDGE), 81100
ATTN TECHNICAL REPORTS BRANCH, 81300
ATTN LEGAL OFFICE, 97000
ATTN ELBAUM, S., 97100
ATTN CHAIRMAN, EDITORIAL COMMITTEE
ATTN ZABLUDOWSKI, B., 47400
ATTN CHIEF, 21000
ATTN CHIEF, 21100
ATTN CHIEF, 21200
ATTN CHIEF, 21300
ATTN CHIEF, 21400
ATTN CHIEF, 21500
ATTN CHIEF, 22000
ATTN CHIEF, 22100
ATTN CHIEF, 22300
ATTN CHIEF, 22800
ATTN CHIEF, 22900
ATTN CHIEF, 20240
ATTN CHIEF, 11000
ATTN CHIEF, 13000
ATTN CHIEF, 13200
ATTN CHIEF, 13300
ATTN CHIEF, 15200
ATTN BROWN, E., 00211
ATTN SINDORIS, A., 00211
ATTN GERLACH, H., 11100
ATTN LIBELO, L., 11200
ATTN LOKERSON, D., 11400
ATTN CROWNE, F., 13200
ATTN DROPKIN, H., 13200
ATTN LEAVITT, R., 13200
ATTN MORRISON, C., 13200
ATTN SATTLER, J., 13200
ATTN KULPA, S., 13300
ATTN SILVERSTEIN, J., 13300
ATTN FAZI, C., 13500
ATTN LOMONACO, S., 15200
ATTN CORRIGAN, J., 20240
ATTN FARRAR, F., 21100
ATTN GARVER, R., 21100
ATTN TATUM, J., 21100
ATTN MERKEL, G., 21300
ATTN MCLEAN, B., 22300
ATTN OLDHAM, T., 22300
ATTN BLACKBURN, J., 22800
ATTN GILBERT, R., 22800
ATTN KLEBERS, J., 22800
ATTN VANDERWALL, J., 22800
ATTN BROMBORSKY, A., 22900

DISTRIBUTION (Cont'd)

ATTN DAVIS, D., 22900
ATTN GRAYBILL, S., 22900
ATTN HUTTLIN, G. A., 22900
ATTN KEHS, A., 22900
ATTN KERRIS, K., 22900
ATTN LAMB, R., 22900
ATTN LINDSAY, D., 22900
ATTN LITZ, M., 22900
ATTN RUTH, B., 22900
ATTN STEWART, A., 22900
ATTN SOLN, J., 22900
ATTN WHITTAKER, D., 22900
ATTN DEHART, J., 47500
ATTN CARRIGAN, J., 20240
ATTN ELBAUM, S., 97100
ATTN BRANDT, H. E., 22300
(60 COPIES)

END

FILMED

7-84

DTIC